
Invasive Rodent Eradication on Islands

GREGG HOWALD,^{*} C. JOSH DONLAN,^{†‡§§§} JUAN PABLO GALVÁN,^{†‡‡‡} JAMES C. RUSSELL,^{§**} JOHN PARKES,^{††} ARACELI SAMANIEGO,^{‡‡} YIWEI WANG,[†] DICK VEITCH,^{§§} PIERO GENOVESI,^{***} MICHEL PASCAL,^{‡‡‡} ALAN SAUNDERS,^{**} AND BERNIE TERSHY[†]

^{*}Island Conservation Canada, 680-220 Cambie Street, Vancouver, British Columbia, V6B 2M9, Canada

[†]Island Conservation, Center for Ocean Health, University of California, 100 Shaffer Road, Santa Cruz, California 95060, U.S.A.

[‡]Department of Ecology and Evolutionary Biology, Corson Hall, Cornell University, Ithaca, New York 14853-2701, U.S.A.

[§]School of Biological Sciences and Department of Statistics, University of Auckland, Private Bag 92019, Auckland, New Zealand, email cjd34@cornell.edu

^{**}Invasive Species Specialist Group, University of Auckland, Tamaki Campus, Private Bag 92019, Auckland, New Zealand

^{††}Landcare Research, P.O. Box 69, Lincoln 8152, New Zealand

^{‡‡}Grupo de Ecología y Conservación de Islas A.C., Avenida López Mateos No. 1590-3, Ensenada, Baja California, C.P. 22880, México

^{§§}48 Manse Road, Papakura 2113, New Zealand

^{***}INFS-Italian Wildlife Institute, Via Ca' Fornacetta 9, Ozzano Emilia (BO) I-40064, Italy

^{‡‡‡}INRA-Equipe Faune Sauvage et Biologie de la Conservation, Station SCRIBE, Campus de Beaulieu, 35 042 Rennes Cedex, France

Abstract: *Invasive mammals are the greatest threat to island biodiversity and invasive rodents are likely responsible for the greatest number of extinctions and ecosystem changes. Techniques for eradicating rodents from islands were developed over 2 decades ago. Since that time there has been a significant development and application of this conservation tool. We reviewed the literature on invasive rodent eradications to assess its current state and identify actions to make it more effective. Worldwide, 332 successful rodent eradications have been undertaken; we identified 35 failed eradications and 20 campaigns of unknown result. Invasive rodents have been eradicated from 284 islands (47,628 ha). With the exception of two small islands, rodenticides were used in all eradication campaigns. Brodifacoum was used in 71% of campaigns and 91% of the total area treated. The most frequent rodenticide distribution methods (from most to least) are bait stations, hand broadcasting, and aerial broadcasting. Nevertheless, campaigns using aerial broadcast made up 76% of the total area treated. Mortality of native vertebrates due to nontarget poisoning has been documented, but affected species quickly recover to pre-eradication population levels or higher. A variety of methods have been developed to mitigate nontarget impacts, and applied research can further aid in minimizing impacts. Land managers should routinely remove invasive rodents from islands <100 ha that lack vertebrates susceptible to nontarget poisoning. For larger islands and those that require nontarget mitigation, expert consultation and greater planning effort are needed. With the exception of house mice (*Mus musculus*), island size may no longer be the limiting factor for rodent eradications; rather, social acceptance and funding may be the main challenges. To be successful, large-scale rodent campaigns should be integrated with programs to improve the livelihoods of residents, island biosecurity, and reinvasion response programs.*

Keywords: eradication, invasive species, island conservation, *Mus musculus*, *Rattus rattus*, *Rattus norvegicus*, *Rattus exulans*

Eradicación de Roedores Invasores de Islas

Resumen: *Los mamíferos invasores son la mayor amenaza a la biodiversidad insular; y los roedores invasores son probables responsables de la mayoría de las extinciones y cambios en los ecosistemas. Las técnicas para la erradicación de roedores de las islas fueron desarrolladas hace 2 décadas. Desde entonces ha habido*

^{‡‡‡}Current address: Sustainable Development and Conservation Biology Program, University of Maryland, College Park, Maryland 20783, U.S.A.

^{§§§}Address correspondence to C.J. Donlan.

Paper submitted November 16, 2006; revised manuscript accepted March 7, 2007.

un desarrollo y aplicación significativa de esta herramienta de conservación. Revisamos la literatura sobre erradicaciones de roedores invasores para evaluar su estado actual e identificar acciones para hacerlo más efectivo. Mundialmente, se han efectuado 332 erradicaciones de roedores exitosas, identificamos 35 erradicaciones fracasadas y 20 campañas con resultados desconocidos. Los roedores Invasivos ha sido erradicados de 284 islas (47,628 ha). Con la excepción de dos islas pequeñas, se utilizaron rodenticidas en todas las erradicaciones. Se utilizó Brodifacoum en 71% de las campañas y en 91% de la superficie tratada. Los métodos más frecuentes de distribución de rodenticida (de más a menos) son estaciones de cebo, aplicación manual y aplicación aérea. Sin embargo, las campañas de aplicación aérea abarcaron 76% de la superficie tratada. Se ha documentado la mortalidad de vertebrados nativos debido a envenenamiento accidental, pero las especies afectadas recuperan, o superan, rápidamente los niveles poblacionales previos a la erradicación. Se ha desarrollado una variedad de métodos para mitigar los impactos no deseados, y la investigación aplicada puede ayudar a minimizar los impactos aun más. Los gestores de recursos deben remover rutinariamente a roedores invasores de islas <100 ha que carezcan de vertebrados susceptibles de envenenamiento no deseado. Para islas más extensas y para las que requieren de mitigación de envenenamientos no deseados, se requiere de la consulta de expertos y de mayores esfuerzos de planificación. Con la excepción de *Mus musculus*, es posible que el tamaño de la isla ya no sea el factor limitante para la erradicación de roedores, más bien, la aceptación social y el financiamiento pueden ser los retos principales. Para ser exitosas, las campañas a gran escala deben estar integradas por programas para mejorar las condiciones de vida de los residentes, de bioseguridad insular y de respuesta a reinvasiones.

Palabras Clave: conservación de islas, erradicación, especies invasoras, *Mus musculus*, *Rattus exulans*, *Rattus norvegicus*, *Rattus rattus*

Introduction

Extinctions over the past thousand years have been dominated by insular species, and invasive mammals have caused the majority of these extinctions (Atkinson 1989; Groombridge et al. 1992). Invasive rodents (rats and house mice [*Mus musculus*]) are likely responsible for the greatest number of extinctions and ecosystem changes on islands (Townsend et al. 2006). Because they are omnivorous, they can affect plants, invertebrates, reptiles, mammals, and birds (Atkinson 1985; Cuthbert & Hilton 2004; Townsend et al. 2006). Invasive rodents occur on over 80% of the world's major islands, and they continue to be introduced onto islands (Atkinson 1985; Pitman et al. 2005).

In response to the negative impacts of invasive rodents on island species and their ecosystems, systematic techniques for eradicating rodents from islands were developed in New Zealand over 2 decades ago (Moors 1985; Taylor & Thomas 1989, 1993). Since then, conservation practitioners have been improving these techniques and leveraging new technologies. As a result, rodents can now be eradicated from larger and biologically complex islands, and eradication has become a powerful tool to prevent extinctions and restore ecosystems (Donlan et al. 2003b; Townsend & Broome 2003). Unfortunately, many invasive rodent eradications remain unpublished or inaccessible, creating the perception among land managers and conservation biologists that successful rodent eradications are rare events (Simberloff 2001; Donlan et al. 2003b). We reviewed invasive rat and house mice eradication campaigns on islands throughout the world. We as-

sessed the approaches, successes, and challenges of these conservation actions to facilitate the conservation of island ecosystems.

Methods

We compiled data from published and gray literature and personal communications on rodent eradications. We judged an eradication campaign a failure or a success based on the outcome reported by the group that conducted the eradication. Because rodents are difficult to detect at low densities (Russell et al. 2005), a widely accepted indicator of eradication success is no detection of rodents after 2 years of intensive monitoring following the eradication effort. Unfortunately, without genetic sampling of rodents on the target island and from potential source populations, it is not possible to distinguish between failure and reinvasion in the first 2–4 years following the eradication effort (Abdelkrim et al. 2007). We did not include secondary eradication efforts of small rodent populations that reinvaded islands following a previous, successful eradication campaign. This is common on islands located close to a mainland source population (Russell & Clout 2005).

All statistical analyses were performed in SPSS, with an α level of 0.05 (SPSS 1999). We used a general linear model to explore relationships of economic costs of eradication campaigns to area, method of baiting, and eradication year. The area covered in an eradication and the cost of the eradication (adjusted to US\$2005) were

\log_{10} transformed to meet normality assumptions. We considered the method of baiting as a categorical fixed effect and modeled the rest of the variables as covariates.

History and Impact of Rodent Introductions

The first rodent (e.g., black rat [*Rattus. rattus*]) introductions to islands may have occurred in the Mediterranean between 5500 and 8000 years ago (Vigne 1992). The kiore (*R. exulans*) was introduced to the islands of the Pacific from Indo-Malaysia some 3000 years ago by the seafaring Lapita people (Atkinson 1985). By approximately 950 years ago, kiore occurred on most of the islands in the Pacific, including New Zealand and likely the Hawaiian and Easter islands (Atkinson 1985; Wilmshurst & Higham 2004). Although exploration by Eurasians may have dispersed black rats to some islands, prior to AD 1500, most islands outside the Pacific were likely free of rats (Atkinson 1985). Between the sixteenth and seventeenth centuries, European explorers spread rats to islands throughout the Indian and Atlantic oceans.

Sometime in the early 1700s, Norway rats (*R. norvegicus*) colonized western Europe, displacing black rats, and subsequently became the dominant species in European and eastern North American ports (Atkinson 1985). Consequently, Norway rats became the dominant rodent on ships and thus the most-introduced rat species on islands throughout the seventeenth and eighteenth centuries. Inexplicably, after the 1850s ship records show that black rats became more common than Norway rats. The presence of both Norway and black rats aboard ships meant that many islands in the Atlantic and Indian oceans had both species and that many Pacific islands had three species.

The distribution of black and Norway rats and kiore on islands worldwide has had devastating effects on island biodiversity. They have negatively affected at least 170 taxa of plants and animals on over 40 islands or archipelagoes and have led to at least 50 extinctions (Townes et al. 2006). Significant indirect and synergistic community- and ecosystem-level effects have also been documented, both in terrestrial and marine environs (Navarrete & Castilla 1993; Imber et al. 2000; Towns 2002; Fukami et al. 2006). Towns et al. (2006) review in detail the biodiversity and ecosystem impacts of *Rattus* spp. in insular environments (for further reference, see Atkinson 1985; Burger & Gochfield 1994).

House mice have had a variety of negative impacts on island ecosystems, including some caused by their predation on reptiles, invertebrates, and the nests of terrestrial birds (Copson 1986; Rowe-Rowe et al. 1989; Newman 1994; Cole et al. 2000; Ruscoe & Murphy 2005). In New Zealand they may have caused the extinction of two invertebrate species on Antipodes Island (Mar-

ris 2000). The effects of house mice on seabird populations are likely underestimated; for example, on Gough Island, house mice prey on Tristan Albatross (*Diomedea dabbenena*) and have significantly reduced the breeding success of colonies (Cuthbert & Hilton 2004). Indirect impacts, such as hyperpredation, of house mice are also probable. They often serve as alternative prey to invasive predators, which in turn can elevate predation levels on native fauna (Bloomer & Bester 1990; Alterio & Moller 1997; Courchamp et al. 2000).

Recovery of insular species following the eradication of invasive rodents is commonplace. Recoveries of terrestrial invertebrates, lizards, and forest birds after eradication have occurred on New Zealand islands (Townes et al. 2006). Seabird populations have responded positively to rat eradications (Jones et al. 2005; Whitworth et al. 2005; Smith et al. 2006; Townes et al. 2006).

Island Rodent Eradications

The first successful rodent eradication was of Norway rats in 1951 on Rouzic Island, France (3.3 ha; Lorvelec & Pascal 2005). Rouzic and early eradications in New Zealand were unintentional byproducts of rodent control efforts (Townes & Broome 2003; see Supplementary Material). Starting in the 1960s and continuing through the mid 1980s, New Zealand conservationists conducted research on bait station approaches and other systematic rodent eradication techniques that resulted in a number of successful intentional eradications on small islands (Moors 1985; Thomas & Taylor 2002). Building on these successes, Norway rats were eradicated from Breaksea Island (170 ha) in 1987, which demonstrated that rodent eradication on larger islands was possible (Taylor & Thomas 1993). The approach used in the Breaksea campaign centered on dispensing a bait containing a proven rodenticide into the territory of every rat with a method that would minimize nontarget poisoning while actively monitoring the progress of the campaign (Taylor & Thomas 1989). Concurrent with the Breaksea and other New Zealand eradication campaigns, black rats were being eradicated on islands in western Australia, including Bodie Island (170 ha; Morris 2002). These research programs and subsequent successful eradications in New Zealand and Australia have spurred hundreds of rodent eradication programs worldwide over the past two decades.

Rodents have been eradicated from at least 284 islands worldwide, totaling over 47,628 ha (Fig. 1; Supplementary Material). Of the known eradication attempts where the result has been documented, 90% have been successful. We documented 387 invasive rodent eradication campaigns, of which 332 were reported successful, 35 failed, and 20 were of unknown outcome. Because successes are more likely to be reported than failures, the success rate may be inflated. On some islands there were




Figure 1. Location and size of islands where successful eradication of invasive rodents have been carried out.

multiple eradication campaigns that either targeted different rodent species or the same species that reinvaded and reestablished after a successful eradication. Most rodent eradications took place in Australasia (155), especially New Zealand (Fig. 1). The majority of rodent eradications have been on islands of <100 ha (78%; Fig. 1). Rats have been removed from 14 islands of over 500 ha. Black rats have been eradicated from most islands worldwide, followed by Norway rats, kiore, and house mice (Table 1). Neither black rats nor house mice have been eradicated from an island larger than 1,000 ha, whereas Norway rats have been removed from Campbell Island, New Zealand, the largest island on which rodent eradication has been successful to date (11,300 ha; Table 1).

Rodenticides

A rodenticide contained in a cereal-based bait was used in all but two small (<14 ha) eradication campaigns (Fig. 2, Supplementary Material). Rodenticide choice and bait depend on a number of factors. The ideal bait is one that is (1) palatable and lethal to the target species after a single feeding event, (2) persistent in the environment long enough for the target species to be exposed

but short enough to minimize nontarget species exposure, (3) has a low probability of engendering bait shyness in target organisms, and (4) is nontoxic or unpalatable to nontarget species. Anticoagulant rodenticides are the most widely used toxin for control of small mammals worldwide (Eason et al. 2002; Hoare & Hare 2006). They act by inhibiting the synthesis of vitamin-K-dependent clotting factors in the liver, which ultimately results in death by internal hemorrhaging, typically within 3–10 days (Hadler & Sahdbolt 1975). Anticoagulants are classified as first- or second-generation according to their potency and when they were developed (Eason et al. 2002). Brodifacoum (3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4 hydroxycoumarin), and other second-generation anticoagulants are more potent with lower LD₅₀ (median lethal dose) values; a single feeding of a few grams of bait can be lethal (Eason et al. 2002). First-generation anticoagulants are less toxic and require multiple feedings over several days to illicit a toxic effect. The higher toxicity and persistence of second-generation anticoagulants is an advantage in eradicating target species; however, that same toxicity and persistence can be a concern when nontarget species are at risk (Hoare & Hare 2006).

Table 1. Invasive rodent eradications: successes, failures, and the largest successful campaign to date.

Species	Successful eradications	Failures (%)	Largest island (ha)*	Method(s)	Reference
<i>Rattus rattus</i>	159	15 (8)	Hermite, AUS (1,022)	aerial broadcast brodifacoum	Burbidge 2004
<i>Rattus norvegicus</i>	104	5 (5)	Campbell, NZL (11,300)	aerial broadcast brodifacoum	McClelland & Tyree 2002
<i>Rattus exulans</i>	55	6 (10)	Hauturu (Little Barrier), NZL (3,083)	aerial broadcast brodifacoum	R. Griffiths, personal communication
<i>Mus musculus</i>	30	7 (19)	Enderby, NZL (710)	aerial broadcast brodifacoum	Torr 2002

*Abbreviations: AUS, Australia; NZL, New Zealand.




Figure 2. Number of successful invasive rodent eradication campaigns by type of rodenticide and method of bait delivery ($n = 264$ islands).

First-generation anticoagulants (i.e., chlorophacinone, diphacinone, pindone, and warfarin) were used in 29 eradication campaigns as the primary rodenticide, and second-generation anticoagulants were used in 226 campaigns (i.e., brodifacoum, bromadiolone, difenacoum, and flocoumafen; Fig. 2). Acute toxins (i.e., 1080 and strychnine) and cholecalciferol were used in six and three campaigns, respectively, as the primary rodenticide (Fig. 2). These nine islands were small (<22 ha), and all but three were supplemented with second-generation anticoagulants (Supplementary Material). Trapping was used to supplement poisoning efforts on 40 islands. Although a number of campaigns used multiple toxins ($n = 33$), this is likely unnecessary unless there are issues with inheritable resistance or high LD_{50} variation (Quy et al. 1995). In 71% of successful campaigns and on 91% of the total area of islands eradicated of invasive rodents, brodifacoum had been applied, making it the most widely used rodenticide.

Bait Delivery

In general the best method for the delivery of a rodenticide depends on island topography, habitat, economics, and vulnerability of nontarget species. The delivery methods currently available are bait stations and hand and aerial broadcasting.

Bait stations, containing rodenticide and distributed on a grid, are the oldest technique used in planned rodent eradication campaigns. Grid sizes vary from 25 to 100 m, depending on the home range of the rodent targeted. Bait stations are monitored and kept filled with rodenticide bait for 1–2 years (Thomas & Taylor 2002). The

bait stations have a number of advantages: they (1) minimize primary exposure to potential nontarget species (e.g., granivorous birds), (2) reduce the amount of toxin delivered to the environment, (3) act as a self-monitoring program with respect to rodenticide uptake, and (4) can be used in combination with nontoxic baits or tracking boards as detection devices after the last rodent supposedly has been killed, which enables managers to kill survivors or immigrants (Thomas & Taylor 2002). Nevertheless, the approach is labor intensive and thus potentially expensive at large scales (e.g., trails might need to be cut), and regular visits to bait stations can result in disturbance of sensitive species, such as breeding seabirds. Furthermore, a bait station approach is impossible with islands that have steep cliffs.

The effectiveness of hand broadcasting was first compared with the bait station technique in 1989 during the eradication of *R. exulans* from Double Island, New Zealand (27 ha). Hand broadcasting proved more cost-effective and led to the development of aerial broadcasting with helicopters (McFadden 1992). Eradication campaigns began using helicopters for aerial broadcast of rodenticides in the early 1990s. Following this, aerial broadcasting was used on larger islands, and hand broadcasting was used on smaller islands (Fig. 3).

Aerial broadcast by helicopter is becoming the most common method of rodenticide delivery (Townes & Broome 2003). Rodenticides can be broadcast on islands with steep and inaccessible cliffs, and aerial or hand broadcasting is often more cost-effective than bait stations. The advent and adoption of geographic positioning systems and geographic information systems technologies have increased the effectiveness and efficiency of




Figure 3. (a) Number of campaigns to eradicate invasive rodents and (b) total area of islands from which rodents have been eradicated with three different methods of bait delivery (percentage of successful campaigns and area of eradication, respectively: bait stations, 54%, 20%; aerial broadcast, 22%, 76%; hand broadcast, 24%, 4% [$n = 269$ islands; large islands > 150 ha]).

invasive mammal eradications, including aerial-based rodent eradications (Lavoie et al. 2007). Because broadcasting entails a single or double bait-application event, usually 10–14 days apart, and bait station campaigns last up to 2 years, broadcasting significantly shortens the eradication campaign (and thus the period of risk to nontarget species). Broadcasting bait in a single application also avoids the issue of cohort selection and interspecific dominance (i.e., where more than one species of target rodent is present), which is likely to arise with the bait stations. In some cases multiple delivery methods may work best. For example, on a small island with steep, accessible cliffs, combining bait stations and hand broadcasting may be the most cost-effective and safest approach. In the end the decision of whether to use bait stations or broadcasting should be one based on experience, consultation, and the constraints of the system (e.g., topography, nontarget species, economics).

The timing of bait delivery also plays a role in eradication planning. Although empirical evidence is scarce (Sweetapple et al. 2002), timing the bait delivery to when

rodents are in decline or during lows in their annual food-dependent population cycle may improve probability of eradication by increasing competition for bait. Timing of bait delivery may also minimize possible nontarget impacts caused by the rodenticide application (e.g., migratory birds) or by the physical nature of the campaign (e.g., disturbing nesting seabirds). With a bait station approach, timing of bait delivery is less of a risk in terms of probability of failure as long as bait remains available throughout food-dependent population declines and long enough for all rodents to gain access to the stations.

The most frequent way of distributing rodenticides was bait stations ($n = 144$) followed by hand broadcasting ($n = 64$) and then aerial broadcasting ($n = 57$; Fig. 2). Nevertheless, aerial broadcast was responsible for 76% of the total area treated. Although bait stations are the most common technique, they have been used on islands of medium size, whereas aerial broadcast has been used on large islands (mean island area of single method campaigns [SE, n], traps = 7.4 ha [$n = 2$], hand broadcast = 20.8 ha [7.3, 37], bait station = 66.2 ha [28.3, 114], aerial

broadcast = 876.4 ha [319.5, 38]). Sixty-seven campaigns used multiple methods (Supplementary Material). Details of hand or aerial broadcast techniques were reported for only 16 campaigns. Of those, rodenticide was delivered in 1–3 applications (mean = 1.56, SE = 0.18, $n = 16$), with a mean application rate of 17.6 kg/ha (median = 15.0, range: 10–35, SE = 2.0, $n = 16$).

Nontarget Species

The risk to nontarget species during an eradication campaign is a function of species present on the island and their behavior; toxicological properties, composition, and delivery method of bait; the susceptibility of those species to the toxin; and the probability of exposure to the toxin either directly by bait consumption or indirectly by feeding on animals that have consumed baits. Although nontarget impacts on vertebrates by primary and secondary poisoning have been documented for eradication campaigns, the affected species have recovered quickly to pre-eradication population levels or higher (Empson & Miskelly 1999; Howald et al. 1999; Davidson & Armstrong 2002; Howald et al. 2005). Invertebrates are less susceptible to anticoagulant toxins. Toxic effects have been elicited in the laboratory, but impacts have not been observed in natural settings, and population-level impacts are unlikely (Booth et al. 2001). Nonetheless, decisions on the choice and delivery of rodenticides, as well as mitigation actions, should be made strategically to minimize any lethal or sublethal impacts on nontarget wildlife (Eason et al. 2002).

Mitigation techniques for vertebrates include live capturing and temporary holding, which has been done successfully for raptors, landbirds, reptiles, and rodents; the use of bait stations in conjunction with an aerial broadcast to provide a selected area as a refuge where rodenticide is not widely available to nontarget species; and the modification of bait stations to limit access to baits by certain species (Townsend et al. 1993; Townsend et al. 1994; Empson & Miskelly 1999; Pergams et al. 2000; Moro 2001; Merton et al. 2002; Morris 2002; Howald et al. 2005). The need to reduce short-term nontarget impacts should be balanced with maximizing the probability of eradication and economic realities (e.g., the lack of funds for a second campaign if an attempt with an alternative toxin fails). Within a holistic framework, a variety of methods are available to mitigate possible nontarget impacts.

Eradication Failures

Eradication failure rates range from 5% for Norway rats to 19% for house mice and depend on the species of rodent, but are only marginally significant ($\chi = 7.32$, $df = 3$, $p = 0.06$, $n = 381$, Table 1). These differences in failure rates highlight the need for more research on house mice eradications, which lag behind in terms of number of successes and largest island successfully targeted. The

cause of these failures is unclear, but they may be related to inadequate bait density in a broadcast application. The home range of house mice is smaller than that of *Rattus*. In general a smaller home range decreases the probability of a target species being exposed to bait that is broadcast at a fixed density over a large area. Additionally, differences in foraging behavior between house mice and *Rattus* could play a role in the dynamics of bait consumption (Macdonald & Fenn 1994).

Managers reported or speculated on causes that contributed to campaign failure in 18 cases (51%). These possible causes included technical issues (e.g., inadequate or insufficient bait deployment), failure to follow established protocols, observed or suspected nontarget poisoning issues that halted the campaign, lack of funding and public support, and bait competition by terrestrial invertebrates.

Economics

We obtained economic costs for only 12% ($n = 47$) of eradication campaigns. Total costs varied widely (US\$123–\$1,615,000, adjusted to 2005 prices), as did cost per hectare (\$3–\$20,000). Not surprisingly, island area and cost of eradication campaign were correlated in log-log space ($F_{1,45} = 76.1$, $p < 0.001$, R^2 [adjusted] = 0.62). A full model, including method of bait delivery (aerial broadcast, hand broadcast, and bait station) and eradication date, did not result in additional significant relationships (method: $F_{3,41} = 0.205$, $p = 0.552$; date: $F_{1,41} = 1.81$, $p = 0.186$; log[area]: $F_{1,41} = 59.9$, $p < 0.001$, R^2 [adjusted] = 0.62). With raw data, area and cost were significantly correlated (Spearman rank correlation: $r_s = 0.746$, $p < 0.01$, $n = 47$).

Martins et al. (2006) claim that eradication costs can be estimated based on limited information, such as area, species, date of eradication, and remoteness. This claim, based on a limited sample size ($n = 41$ for all invasive mammals), is disconnected from the many realities of the costs of eradications (Donlan & Wilcox 2007). In addition to area, remoteness, and target species, the costs of eradication campaigns can differ drastically depending on a suite of fixed and nonfixed costs, including mitigation for potential nontarget species, techniques used, local capacity and bureaucracy, and the environmental compliance required (Donlan & Wilcox 2007).

Challenges and Recommendations

The eradication of invasive rodents from islands, like other invasive mammals, is no longer a rare event (Nogales et al. 2004; Campbell & Donlan 2005). Rather, it is a powerful tool to prevent further extinctions and to restore ecosystems (Hutton et al. 2007), often with high conservation returns from a cost-benefit perspective. For

example, 201 seabird colonies and 88 endemic terrestrial vertebrates have been protected on the islands of western Mexico through invasive mammal eradication at a cost of US\$21,000 and US\$49,000 per colony or taxon, respectively (Aguirre-Muñoz et al. 2007). In addition to negative biodiversity impacts, rodents also affect people living on islands through their degradation of food crops and their role as disease vectors (Hood et al. 1971; Chanteau et al. 1998). Thus, rodent eradication can also result in social and economic benefits. For example, the residents of Lord Howe Island, Australia, have proposed eradicating rodents to reduce the economic impacts on agriculture (A.S., personal observation).

With proper preparation (Cromarty et al. 2002), land managers should routinely remove invasive rodents from islands <100 ha that lack native vertebrates susceptible to nontarget poisoning. For larger islands and islands with potential nontarget poisoning issues, land managers should seek expert consultation from experienced practitioners. Additional planning focused on the type, timing, and delivery method of rodenticide and on mitigating potential nontarget impacts is needed. Whenever possible the negative effects of the eradication process and the benefits of the island being rodent-free should be documented in a monitoring program. Furthermore, invasive mammal eradication offers unique opportunities for large-scale ecological experiments (Donlan et al. 2002; Croll et al. 2005). At the least, eradication campaigns should report success or failure and economic costs. A public database is available for reporting on eradication campaigns (<http://www.issg.org>). Development of global and regional prioritization models to elucidate where to invest in rodents and other invasive species eradication to maximize biodiversity gained on the investment should be a high priority.

Eradication campaigns can face opposition from individuals or organizations concerned about animal rights or toxicity issues (Towns et al. 2006). For example, on Anacapa Island (California, U.S.A.), an animal rights organization filed an unsuccessful legal injunction to halt a rat eradication (Howald et al. 2005). As larger islands, many of them with human populations, are targeted for eradication, incorporating human dimensions into eradication planning will be increasingly important (Genovesi 2007).

Conservationists must also work with regulatory agencies on a nuanced set of laws that protect people and wildlife in continental settings, but maintain the use of a suite of useful rodenticides. For example, in the United States and United Kingdom there are serious concerns and issues with nontarget rodenticide poisoning of birds and mammals due to the widespread availability, chronic use, and misuse of brodifacoum. These concerns have resulted in calls for wholesale restrictions (Stone et al. 1999; Fournier-Chambrillon et al. 2004; Brakes & Smith 2005). This level of use of brodifacoum is vastly different than a one-time, restricted rodenticide application on an island

for conservation purposes. Although increased regulation on certain rodenticides may be justified, brodifacoum is currently the most important rodenticide for invasive rodent eradication on islands and should remain available to practitioners to use responsibly.

Applied research can help eradication campaigns minimize potential nontarget impacts of native wildlife while maximizing probability of eradication success. Collaborative research is underway to explore the possibilities of a toxin specific to *Rattus*. Invasive and native rodents are equally susceptible to available rodenticides. To date, invasive rodents have been eradicated from only two islands with an endemic terrestrial mammal (Morris 1989; Howald et al. 2005). Both *Rattus*- and *Mus*-specific toxins would have substantial global conservation implications, particularly on islands with endemic terrestrial rodents and endemic birds susceptible to nontarget poisoning. Research is also needed to test the field efficacy of alternative toxins and lower application rates that could minimize potential nontarget impacts and reduce the amount of toxin released into the environment. Small islands are the ideal testing grounds for this research. Encouragingly, diphacinone and cholecalciferol, which are less toxic to birds, have been used successfully in four rodent campaigns on small islands (Donlan et al. 2003a; Smith et al. 2006; Witmer et al. 2007). Finally, more research is needed on house mice eradication and invasive rodent eradication in tropical environments, where bait competition with terrestrial invertebrates (e.g., land crabs) presents unique challenges (Rodríguez et al. 2006).

A significant risk that has yet to be addressed adequately in aerial baiting strategies is the inability to detect, locate, and address potential survivors of eradication campaigns. Current practice is to plan carefully and hope the campaign kills 100% of the rodents. Failure is assessed by waiting until such time as survivors could have produced enough offspring for the population to become easily detectable. This approach assumes that it would cost more to detect and locate potential survivors than to repeat the entire eradication campaign. Tactical research is needed to shift this cost-benefit differential toward timely posteradication detection (e.g., highly trained dogs) and response. Such response is required for rabbit eradication, where 100% of the population is never killed during initial aerial baiting campaigns and for other species for which eradication is achieved via repeated harvesting (Parkes 2006). Additionally, managers need decision tools to determine when it is cost-effective to switch management schemes from active eradication to monitoring (Cacho et al. 2006; Regan et al. 2006).

Conservation practitioners are now eradicating invasive rodents from larger and more biologically complex islands. As larger islands are targeted, a number of factors will become increasingly important: rodenticide choice and the development of new rodenticides, minimizing nontarget and secondary poisoning events, and

leveraging technology to allow techniques to scale from smaller to larger islands. With the exception of house mice, island size may no longer be the most limiting factor with respect to the ability to remove invasive rodents; rather, nontarget impacts, sociology, and funding will be the main challenges. Because of the presence of humans on many larger islands, future rodent eradication will require integrated environmental education, island biosecurity, and reinvasion response programs. Failure to maintain adequate island biosecurity regimes can lead to reinvasions, which can be difficult to detect and to mount a response against. Increasing the efficiency of eradication, including bioeconomic analyses, will also be important because absolute costs, probability of failure, and conservation benefits will all increase with the size of the island (e.g., Choquenot & Parkes 2001; Choquenot 2006). A large percentage of the world's threatened biodiversity resides on islands where invasive species are the major threat. Because it is possible to safely eradicate invasive rodents from islands and because there is a high return in biodiversity gains following eradication, invasive rodents should be routinely removed from islands.

Acknowledgments

We thank the many people who provided data, contacts, internal reports, and other information: M. Abrose, N. Baccetti, E. Bell, M. Browne, A. Burbidge, K. Campbell, V. Carrion, F. Courchamp, J. Daltry, M. Evans, J. Fric, G. Gerber, R. Griffiths, J. Hughes, K. Lindsay, J. Mackenzie, I. McFadden, G. Meier, D. Merton, R. Parrish, J. Ramdirez, B. Simmons, P. Sposimo, K. Swift, D. P. Taylor, P. Thomson, K. Varnham, G. Wragg, B. Zonfrillo. C.J.D. thanks J. Berger and K. M. Berger for sharing their space and K. A. Berger and Z. B. Berger for balance. Funding was provided by Cornell University to C.J.D.

Supplementary Material

Characteristics of islands where invasive rodent eradication failed or was successful or of unknown outcome (Appendix S1) are available as part of the on-line article from <http://www.blackwell-synergy.com/>. The author is responsible for the content and functionality of these materials. Queries (other than absence of the material) should be directed to the corresponding author.

Literature Cited

- Abdelkrim, J., M. Pascal, and S. Samadi. 2007. Establishing causes of eradication failure based on genetics: case study of ship rat eradication in Ste. Anne archipelago. *Conservation Biology* **21**:719–730.
- Alterio, N. K., and H. Moller. 1997. Diet of feral house cats *Felis catus*, ferrets *Mustela furo*, and stoats *M. erminea* in grassland surrounding yellow-eyed penguin *Megadyptes antipodes* breeding areas, South Island, New Zealand. *Journal of Zoology* **243**:869–877.
- Aguirre-Muñoz, A., et al. 2007. High-impact conservation action: a case study from the islands of western Mexico. *Ambio*: in press.
- Atkinson, I. A. E. 1985. The spread of commensal species of *Rattus* to oceanic islands and their effects on island avifaunas. Pages 35–81 in P. J. Moors, editor. *Conservation of island birds: case studies for the management of threatened island species*. International Council for Bird Preservation, Cambridge, United Kingdom.
- Atkinson, I. A. E. 1989. Introduced animals and extinctions. Pages 54–75 in D. Western and M. C. Pearl, editors. *Conservation for the twenty-first century*. Oxford University Press, New York.
- Bloomer, J. P., and M. N. Bester. 1990. Diet of a declining feral cat *Felis catus* population on Marion Island (Antarctic). *South African Journal of Wildlife Research* **20**:1–4.
- Booth, L. H., C. T. Eason, and E. B. Spurr. 2001. Literature review of the acute toxicity and persistence of brodifacoum to invertebrates. *Science for Conservation* **177**:1–9.
- Brakes, C. R., and R. H. Smith. 2005. Exposure of nontarget small mammals to rodenticides: short-term effects, recovery, and implications for secondary poisoning. *Journal of Applied Ecology* **42**:118–128.
- Burbidge, A. A. 2004. Montebello renewal: Western Shield review—February 2003. *Conservation Science Western Australia* **5**:194–201.
- Burger, J., and M. Gochfield. 1994. Predation and effects of humans on island nesting birds. Pages 39–67 in D. N. Nettleship, J. Burger, and M. Gochfield, editors. *Seabirds on islands, threats, case studies, and action plans*. Birdlife Conservation Series, Hamilton, New Zealand.
- Cacho, J. O., D. Spring, P. Pheloung, and S. Hester. 2006. Evaluating the feasibility of eradicating an invasion. *Biological Invasions* **8**:903–917.
- Campbell, K., and C. J. Donlan. 2005. Feral goat eradication on islands. *Conservation Biology* **19**:1362–1374.
- Chanteau, S., L. Ratsifasoamanana, B. Rasoamanana, L. Rahalison, J. Randriambeloso, J. Roux, and D. Rabeson. 1998. Plague, a reemerging disease in Madagascar. *Emerging Infectious Diseases* **4**:101–104.
- Choquenot, D. 2006. Bioeconomic modeling in conservation pest management: effect of stoat control on extinction risk of an indigenous New Zealand Passerine, *Moboua ochrocephala*. *Conservation Biology* **20**:480–489.
- Choquenot, D., and J. Parkes. 2001. Setting thresholds for pest control: how does pest density affect resource viability? *Biological Conservation* **99**:29–46.
- Cole, F. R., L. L. Loope, A. C. Medeiros, C. E. Howe, and L. J. Anderson. 2000. Food habits of introduced rodents in high-elevation shrubland of Haleakala National Park, Maui, Hawai'i. *Pacific Science* **54**:313–329.
- Copson, G. R. 1986. The diet of the introduced rodents *Mus musculus* and *Rattus rattus* on subantarctic Macquarie Island. *Australian Wildlife Research* **13**:441–446.
- Courchamp, F., M. Langlais, and G. Sugihara. 2000. Rabbits killing birds: modeling the hyperpredation process. *Journal of Animal Ecology* **69**:154–164.
- Croll, D. A., J. L. Maron, J. A. Estes, E. M. Danner, and G. V. Byrd. 2005. Introduced predators transform subarctic islands from grassland to tundra. *Science* **307**:1959–1961.
- Cromarty, P. L., K. G. Broome, A. Cox, R. A. Empson, W. M. Hutchinson, and I. McFadden. 2002. Eradication planning for invasive alien animal species on islands—the approach developed by the New Zealand Department of Conservation. Pages 85–91 in C. R. Veitch and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. World Conservation Union, SSC Invasive Species Specialist Group, Gland, Switzerland.
- Cuthbert, R., and G. Hilton. 2004. Introduced house mice *Mus musculus*: a significant predator of threatened and endemic birds on Gough Island, South Atlantic Ocean? *Biological Conservation* **117**:483–489.

- Davidson, R. S., and D. P. Armstrong. 2002. Estimating impacts of poison operations on nontarget species using mark-recapture analysis and simulation modeling: an example with saddlebacks. *Biological Conservation* **105**:375–381.
- Donlan, C. J., B. R. Tershy, and D. A. Croll. 2002. Islands and introduced herbivores: conservation action as ecosystem experimentation. *Journal of Applied Ecology* **39**:235–246.
- Donlan, C. J., G. R. Howald, B. R. Tershy, and D. A. Croll. 2003a. Evaluating alternative rodenticides for island conservation: roof rat eradication from the San Jorge Islands, Mexico. *Biological Conservation* **114**:29–34.
- Donlan, C. J., B. R. Tershy, K. Campbell, and F. Cruz. 2003b. Research for requiems: the need for more collaborative action in invasive species management and conservation. *Conservation Biology* **17**:1850–1851.
- Donlan, C. J., and C. Wilcox. 2007. The complexities of costing eradications. *Animal Conservation* **10**:156–158.
- Eason, C. T., E. C. Murphy, G. R. G. Wright, and E. B. Spurr. 2002. Assessment of risks to brodifacoum to non-target birds and mammals to New Zealand. *Ecotoxicology* **11**:35–48.
- Empson, R. A., and C. M. Miskelly. 1999. The risks, costs and benefits of using brodifacoum to eradicate rats from Kapiti Island, New Zealand. *New Zealand Journal of Ecology* **23**:241–254.
- Fournier-Chambrillon, C., et al. 2004. Evidence of secondary poisoning of free-ranging riparian mustelids by anticoagulant rodenticides in France: implications for conservation of European mink (*Mustela lutreola*). *Journal of Wildlife Diseases* **40**:688–695.
- Fukami, T., D. A. Wardle, P. J. Bellingham, C. P. H. Mulder, D. R. Towns, G. W. Yeates, K. I. Bonner, M. S. Durrett, M. N. Grant-Hoffman, and W. M. Williamson. 2006. Above- and below-ground impacts of introduced predators in seabird-dominated island ecosystems. *Ecology Letters* **9**:1299–1307.
- Genovesi, P. 2007. Limits and potentialities of eradication as a tool for addressing biological invasions. Pages 385–400 in W. Nentwig, editor. *Biological invasions. Ecological studies*. Volume 193. Springer, New York.
- Groombridge, B. 1992. World Conservation Monitoring Centre, British Natural History Museum, and World Conservation Union. *Global biodiversity: status of the earth's living resources: a report*. Chapman & Hall, London.
- Hadler, M. R., and R. S. Sahdbolt. 1975. Novel 4-hydroxy-coumarin anticoagulants active against resident rats. *Nature* **253**:277–282.
- Hoare, J. M., and K. M. Hare. 2006. The impact of brodifacoum on nontarget wildlife: gaps in knowledge. *New Zealand Journal of Ecology* **30**:157–167.
- Hood, G. A., R. D. Nass, G. D. Lindsey, and D. N. Hirata. 1971. Distribution and accumulation of rat damage in Hawaiian sugarcane. *Journal of Wildlife Management* **35**:613–618.
- Howald, G. R., P. Mineau, J. E. Elliott, and K. M. Cheng. 1999. Brodifacoum poisoning of avian scavengers during rat control on a seabird colony. *Ecotoxicology* **8**:431–447.
- Howald, G. R., K. R. Faulkner, B. Tershy, B. Keitt, H. Gellerman, E. M. Creel, M. Grinnel, S. Ortega, and D. A. Croll. 2005. Eradication of black rat from Anacapa Island: biological and social considerations. Pages 299–312 in D. K. Garcelon and C. A. Schwemm, editors. *Proceedings of the sixth California Islands symposium*. Institute for Wildlife Studies, Arcata, California.
- Hutton, I., J. P. Parkes, and A. R. E. Sinclair. 2007. Reassembling island ecosystems: the case of Lord Howe Island. *Animal Conservation* **10**:22–29.
- Imber, M., M. Harrison, and J. Harrison. 2000. Interactions between petrels, rats and rabbits on Whale Island, and effects of rat and rabbit eradication. *New Zealand Journal of Ecology* **24**:153–160.
- Jones, H. P., R. W. Henry, G. R. Howald, B. R. Tershy, and D. A. Croll. 2005. Predation of artificial Xantus's Murrelet (*Synthliboramphus hypoleucus scrippsi*) nests before and after black rat (*Rattus rattus*) eradication. *Environmental Conservation* **32**:320–325.
- Lavoie, C., C. J. Donlan, K. Campbell, F. Cruz, and V. Carrion G. 2007. Geographic tools for eradication programs of insular non-native mammals. *Biological Invasions* **9**:139–148.
- Lorvelec, O., and M. Pascal. 2005. French attempts to eradicate non-indigenous mammals and their consequences for native biota. *Biological Invasions* **7**:135–140.
- Macdonald, D. W., and M. G. P. Fenn. 1994. The natural history of rodents. Pages 1–22 in A. P. Buckle and R. H. Fenn, editors. *Rodent pests and their control*. CAB International, Wallingford, United Kingdom.
- Marris, J. W. M. 2000. The beetle (Coleoptera) fauna of the Antipodes Islands, with comments on the impact of mice: and an annotated checklist of the insect and arachnid fauna. *Journal of the Royal Society of New Zealand* **30**:169–195.
- Martins, T. L. E., M. D. L. Brooke, G. M. Hilton, S. Farnsworth, J. Gould, and D. J. Pain. 2006. Costing eradication of alien mammals from islands. *Animal Conservation* **9**:439–444.
- McClelland, P., and P. Tyree. 2002. Eradication: the clearance of Campbell Island. *New Zealand Geographic* **58**:86–94.
- McFadden, I. 1992. Eradication of kiore (*Rattus exulans*) from Double Island, Mercury Group, in northern New Zealand. Internal report 130. Department of Conservation, Science and Research, Wellington, New Zealand.
- Merton, D., G. Climo, V. Laboudallon, S. Robert, and C. Mander. 2002. Alien mammal eradication and quarantine on inhabited islands in the Seychelles. Pages 182–198 in C. R. Veitch and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. World Conservation Union, SSC Invasive Species Specialist Group, Gland, Switzerland.
- Moors, P. J. 1985. Eradication campaign against *Rattus norvegicus* on the Noises Islands, New Zealand, using brodifacoum and 1080. International Council for Bird Preservation, Technical publication **3**:143–155.
- Moro, D. 2001. Evaluation and cost benefits of controlling house mice *Mus domesticus* on islands: an example from Thevenard Island, Western Australia. *Biological Conservation* **99**:355–664.
- Morris, K. D. 1989. Feral animal control on Western Australian islands. Pages 105–111 in A. Burbidge, editor. *Australian and New Zealand islands: nature conservation values and management*. Department of Conservation and Land Management, Perth.
- Morris, K. D. 2002. The eradication of the black rat (*Rattus rattus*) on Barrow and adjacent islands off the northwest coast of Western Australia. Pages 219–225 in C. R. Veitch and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. World Conservation Union, SSC Invasive Species Specialist Group, Gland, Switzerland.
- Navarrete, S. A., and J. C. Castilla. 1993. Predation by Norway rats in the intertidal zone of central Chile. *Marine Ecology Progress Series* **92**:187–199.
- Newman, D. G. 1994. Effects of a mouse, *Mus musculus*, eradication programme and habitat change on a lizard populations on Mana Island, New Zealand, with special reference to McGregor's skink, *Cyclodina macgregori*. *New Zealand Journal of Zoology* **21**:443–456.
- Nogales, M., A. Martín, B. R. Tershy, C. J. Donlan, D. Veitch, N. Puerta, B. Wood, and J. Alonso. 2004. A review of feral cat eradication on islands. *Conservation Biology* **18**:310–319.
- Parkes, J. 2006. Eradication of vertebrate pests: are there any general lessons? Pages 91–110 in C. J. Feare and D. P. Cowan, editors. *Advances in vertebrate pest management IV*. Filander Verlag, Furth, Germany.
- Pergams, O. R. W., R. C. Lacy, and M. V. Ashley. 2000. Conservation and management of Anacapa Island deer mice. *Conservation Biology* **14**:819–832.
- Pitman, R. L., L. T. Ballance, and C. Bost. 2005. Clipperton island: pig sty, rat hole and booby prize. *Marine Ornithology* **33**:193–194.
- Quy, R. J., D. P. Cowan, C. V. Prescott, J. E. Gill, G. M. Kerrins, G. Dunsford, A. Jones, and A. D. McNicholl. 1995. Control of a population of Norway rats resistant to anticoagulant rodenticides. *Pesticide Science* **45**:247–256.

- Regan, T. J., M. A. McCarthy, P. W. J. Baxter, F. D. Panetta, and H. P. Possingham. 2006. Optimal eradication: when to stop looking for an invasive plant. *Ecology Letters* 9:759–766.
- Rodríguez, C., R. Torres, and H. Drummond. 2006. Eradicating introduced mammals from a forested tropical island. *Biological Conservation* 130:98–105.
- Rowe-Rowe, D. T., B. Green, and J. E. Crafford. 1989. Estimated impact of feral house mice on subantarctic invertebrates at Marion Island [Indian Ocean]. *Polar Biology* 9:457–460.
- Ruscoe, W. A., and E. C. Murphy. 2005. House mouse. Pages 204–221 in C. M. King, editor. *The handbook of New Zealand mammals*. Oxford University Press, Melbourne.
- Russell, J. C., and M. N. Clout. 2005. Rodent incursions on New Zealand islands. Pages 324–330 in J. Parkes, M. Statham, and G. Edwards, editors. *Proceedings of the 13th Australasian vertebrate pest conference*. Landcare Research, Lincoln, New Zealand.
- Russell, J. C., D. R. Towns, S. H. Anderson, and M. N. Clout. 2005. Intercepting the first rat ashore. *Nature* 437:7062.
- Simberloff, D. 2001. Eradication of island invasives: practical actions and results achieved. *Trends in Ecology & Evolution* 16:273–274.
- Smith, D. G., E. K. Shiinoki, and E. A. VanderWerf. 2006. Recovery of native species following rat eradication on Mokoli'i, O'ahu, Hawai'i. *Pacific Science* 60:299–303.
- SPSS. 1999. SPSS Base 10.0 for Window user's guide. SPSS, Chicago.
- Stone, W. B., J. C. Okonewski, and J. R. Stedelin. 1999. Poisoning of wildlife with anticoagulant rodenticides in New York. *Journal of Wildlife Diseases* 35:187–193.
- Sweetapple, P., G. Nugent, and W. Fraser. 2002. Long-term changes in rat abundance following possum control. *Possum Research News* 17:1–2.
- Taylor, R. H., and B. W. Thomas. 1989. Eradication of Norway rats *Rattus norvegicus* from Hawea Island, Fiordland, using brodifacoum. *New Zealand Journal of Ecology* 12:23–32.
- Taylor, R. H., and B. W. Thomas. 1993. Rats eradicated from rugged Breaksea island (170 Ha), Fiordland, New Zealand. *Biological Conservation* 65:191–198.
- Thomas, B. W., and R. H. Taylor. 2002. A history of ground-based rodent eradication techniques developed in New Zealand, 1959–1993. Pages 301–310 in C. R. Veitch and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. World Conservation Union, SSC Invasive Species Specialist Group, Gland, Switzerland.
- Torr, N. 2002. Eradication of rabbits and mice from sub-Antarctic Enderby and Rose islands. Pages 319–328 in C. R. Veitch and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. World Conservation Union, SSC Invasive Species Specialist Group, Gland, Switzerland.
- Towns, D. R. 2002. Interactions between geckoes, honeydew scale insects and host plants revealed on islands in northern New Zealand, following eradication of introduced rats and rabbits. Pages 329–335 in C. R. Veitch and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. World Conservation Union, SSC Invasive Species Specialist Group, Gland, Switzerland.
- Towns, D. R., and K. G. Broome. 2003. From small Maria to massive Campbell: forty years of rat eradications from New Zealand islands. *New Zealand Journal of Zoology* 30:377–398.
- Towns, D. R., I. McFadden, and T. Lovegrove. 1993. Offshore islands co-operative conservation project with ICI Crop Care Division: phase one (Stanley Island). Internal report 138. Department of Conservation, Science and Research Wellington, New Zealand.
- Towns, D. R., M. I., P. Thomson, H. Robertson, and R. Colbourne. 1994. Offshore islands co-operative conservation project with ICI Crop Care Division: phase two (Red Mercury Island). Internal report 142. Department of Conservation, Science and Research, Wellington, New Zealand.
- Towns, D. R., I. A. E. Atkinson, and C. H. Daugherty. 2006. Have the harmful effects of introduced rats on islands been exaggerated? *Biological Invasions* 8:863–891.
- Vigne, J. D. 1992. Zooarchaeology and the biogeographical history of mammals of Corsica and Sardinia since the last ice age. *Mammal Review* 22:87–96.
- Whitworth, D. L., H. R. Carter, R. J. Young, J. S. Koepke, F. Gress, and S. Fangman. 2005. Initial recovery of Xantus's Murrelets following rat eradication on Anacapa Island, California. *Marine Ornithology* 33:131–137.
- Wilmschurst, J., and T. E. G. Higham. 2004. Using rat-gnawed seeds to independently date the arrival of Pacific rats and humans in New Zealand. *Holocene* 14:801–806.
- Witmer, G. W., F. Boyd, and Z. Hillis-Star. 2007. The successful eradication of introduced roof rats (*Rattus rattus*) on Buck Island using diphacinone, followed by resource recovery and an irruption of house mice (*Mus musculus*). *Wildlife Research* 34: 108–115.

Appendix 1. Characteristics of islands where invasive rodents have been successfully eradicated, failed, or of unknown outcome. Primary (1°), secondary (2°) and tertiary (3°) methods and rodenticide are included.

SUCCESSFUL ERADICATIONS

<i>Island (Alternative Name), Group</i>	<i>County^a</i>	<i>Species</i>	<i>Area (ha)</i>	<i>Year Eradicated</i>	<i>Methods (1°,2°,3°)</i>	<i>Rodenticide (1°,2°,3°)</i>	<i>Reference</i>
Galley Islands (2 islets), North Sound	A&B	R. rattus	1	1995	bait stations	brodifacoum	J. Daltry pers. comm.
Lobster, North Sound	A&B	R. rattus	1	1998	bait stations	brodifacoum	K. Varnham pers. comm
Redhead, North Sound	A&B	R. rattus	3	1998	bait stations	brodifacoum	K. Varnham pers. comm.
Rabbit, North Sound	A&B	R. rattus	6	1998	bait stations	brodifacoum	K. Varnham pers. comm
Great Bird, North Sound	A&B	R. rattus	10	1995	bait stations	brodifacoum	K. Lindsay pers. comm.
Great Bird, North Sound	A&B	R. rattus	10	2002	bait stations	brodifacoum	K. Lindsay pers. comm.
Green, North Sound	A&B	R. rattus	43	2001	bait stations	brodifacoum	K. Varnham pers. comm
Beacon, Lowendal	AUS	M. musculus	1.2	1997	bait stations	pindone, brodifacoum	Burbidge & Morris 2002
Bridled, Lowendal	AUS	M. musculus	22	1997	bait stations	pindone, brodifacoum	Burbidge & Morris 2002
Varanus, Lowendal	AUS	M. musculus	80	1997	bait stations	pindone, brodifacoum	Burbidge & Morris 2002
Barrow ^b	AUS	M. musculus		1965	bait stations	pindone	Burbidge & Morris 2002
Barrow ^b	AUS	M. musculus		1972	bait stations	pindone	Burbidge & Morris 2002
Barrow ^b	AUS	M. musculus		1994	bait stations	pindone	Burbidge & Morris 2002
Barrow ^b	AUS	M. musculus		1998	bait stations	pindone	Burbidge & Morris 2002
Fisher, Furneaux	AUS	R. rattus	1				Abbott & Burbidge 1995
Pasco, Barrow	AUS	R. rattus	2	1985	bait stations	pindone	Morris 1989; Morris 2002
Pigeon, Houtman Albrolhos	AUS	R. rattus	3	1993	hand broadcast	brodifacoum	Burbidge & Morris 2002; A. Burbidge unpublished data
Bloodwood, Montebellos	AUS	R. rattus	3	1996	bait stations	brodifacoum	Burbidge 2004
Ivy, Montebellos	AUS	R. rattus	4	1996	bait stations	brodifacoum	Burbidge 2004

Prince, Barrow	AUS	R. rattus	4	1983	bait stations	pindone	Morris 1989, Burbidge & Morris 2002
Boomerang, Barrow	AUS	R. rattus	5	1983	bait stations	pindone	Morris 1989; Morris 2002
Sandy, Lacepede	AUS	R. rattus	6	1986	bait stations	pindone	Burbidge & Morris 2002
Double North, Barrow	AUS	R. rattus	12	1983	bait stations	pindone	Morris 1989; Morris 2002
Heron, Capricorn	AUS	R. rattus	13				Abbott & Burbidge 1995
South East, Montebellos	AUS	R. rattus	13	1996	bait stations	brodifacoum	Burbidge 2004
Brooke, Montebellos	AUS	R. rattus	15	1996	bait stations	brodifacoum	Burbidge 2004
Ah Chong, Montebellos	AUS	R. rattus	22	1996	bait stations	brodifacoum	Burbidge 2004
Double South, Barrow	AUS	R. rattus	23	1983	bait stations	pindone	Morris 1989; Morris 2002
Bedout	AUS	R. rattus	24	1981	bait stations	pindone	Morris 1989; Burbidge & Morris 2002
Delta, Montebellos	AUS	R. rattus	38	1999	bait stations	brodifacoum	Burbidge 2004
Crocus, Montebellos	AUS	R. rattus	41	1997	bait stations	brodifacoum	Burbidge 2004
Primose, Montebellos	AUS	R. rattus	41	1997	bait stations	brodifacoum	Burbidge 2004
Campbell, Montebellos	AUS	R. rattus	47	1999	bait stations	brodifacoum	Burbidge 2004
Rat, Houtman Abrolhos	AUS	R. rattus	56	1993	bait stations	brodifacoum	Burbidge & Morris 2002; A. Burbidge unpublished data
Renewal, Montebellos	AUS	R. rattus	58	1996	bait stations	brodifacoum	Burbidge 2004
Middle, Lacepede	AUS	R. rattus	60	1986	bait stations	pindone	Burbidge & Morris 2002
Bluebell, Montebellos	AUS	R. rattus	65	1996	bait stations	brodifacoum	Burbidge 2004
Bluebell, Montebellos	AUS	R. rattus	65	2001	aerial broadcast	brodifacoum	Burbidge 2004
West, Lacepede	AUS	R. rattus	82	1986	bait stations	pindone	Burbidge & Morris 2002
Alpha, Montebellos	AUS	R. rattus	118	1996	bait stations	brodifacoum	Burbidge 2004
Alpha, Montebellos	AUS	R. rattus	118	2001	aerial broadcast	brodifacoum	Burbidge 2004
Northwest, Montebellos	AUS	R. rattus	135	1996	bait stations	brodifacoum	Burbidge 2004
Boodie, Barrow	AUS	R. rattus	170	1985	bait stations	pindone	Morris 1989; Morris 2002
Middle, Barrow	AUS	R. rattus	350	1991	bait stations	pindone	Morris 2002, Burbidge & Morris 2002
Trimouille, Montebellos	AUS	R. rattus	522	1996	bait stations	brodifacoum	Burbidge 2004
Hermite, Montebellos	AUS	R. rattus	1022	2001	aerial broadcast	brodifacoum	Burbidge 2004
Barrow	AUS	R. rattus	270 (of 23,000)	1991	bait stations	pindone	Morris 2002
Panay, Montebello	AUS	R. rattus		1996	bait stations	brodifacoum	Burbidge 2004
White Cay (Sandy Cay), Exumas	BAH	M. musculus	15	1998	bait stations	brodifacoum	Hayes et al. 2004
Low Cay, San Salvador	BAH	R. rattus	10.8	2000	bait stations	brodifacoum	Hayes et al. 2004
White Cay (Sandy Cay), Exumas	BAH	R. rattus	15	1998	bait stations	brodifacoum	Hayes et al. 2004

Cox, Queen Charlottes	CAN	R. norvegicus	10	1995	bait stations	brodifacoum	Kaiser et al. 1997
Lucy, Queen Charlottes	CAN	R. norvegicus	40		bait stations	brodifacoum	Buck 1995
Langara, Queen Charlottes	CAN	R. norvegicus	3105	1995	bait stations	brodifacoum	Kaiser et al. 1997
Anchorage, Suvarrow	COI	R. exulans	12	2003	hand broadcast	brodifacoum	G. Wragg pers. comm
Pitt, Galapagos	ECU	R. rattus	0.4	1989	hand broadcast, traps	1080	K. Campbell pers. comm.
Bainbridge-2, Galapagos	ECU	R. rattus	2.9	2002	hand broadcast	brodifacoum	K. Campbell pers. comm.
Mosquera, Galapagos	ECU	R. rattus	4.6		hand broadcast	brodifacoum, 1080	K. Campbell pers. comm.
Lobos, Galapagos	ECU	R. rattus	6.7	2002	bait stations, traps	brodifacoum	V. Carrion pers. comm
Surprise, New Caledonia	FRA	M. musculus	24	2005	hand broadcast	bromadiolone	F. Courchamp pers. comm.
Fajou, Guadeloupe	FRA	M. musculus	120	2001	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005, M. Pascal unpublished data
Laregnere, New Caledonia	FRA	R. exulans	0.5	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Ndo, New Caledonia	FRA	R. exulans	2	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Uie, New Caledonia	FRA	R. exulans	2	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Uo, New Caledonia	FRA	R. exulans	3	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Nge, New Caledonia	FRA	R. exulans	4	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
G'i, New Caledonia	FRA	R. exulans	5	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Uatermbi, New Caledonia	FRA	R. exulans	5	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Uatio, New Caledonia	FRA	R. exulans	5	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Vua, New Caledonia	FRA	R. exulans	5	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Redika, New Caledonia	FRA	R. exulans	7	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Rocher de Cancale, Rimains	FRA	R. norvegicus	0.2	1994	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005, Abdelkrim et al. 2005a
Île aux Rats, Sept-Île	FRA	R. norvegicus	0.2	1994	bait stations, traps	bromadiolone	Lorvelec & Pascal 2005, Pascal et al. 2005
Chatellier, Rimains	FRA	R. norvegicus	1	1994	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005, Abdelkrim et al. 2005b
Enez ar C'hrizienn, Molène	FRA	R. norvegicus	1.3	1996	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005

Rimains, Rimains	FRA	R. norvegicus	1.5	1994	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005, Abdelkrim et al. 2005a
Île aux Chevaux , Houat	FRA	R. norvegicus	2.5	2002	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005
Rouzic ,Sept-Île	FRA	R. norvegicus	3.3	1951	hand broadcast	strychnine	Lorvelec & Pascal 2005
Île Plate, Sept-Île	FRA	R. norvegicus	5	1994	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005, Pascal et al. 2005
Île aux Moines, Sept-Île	FRA	R. norvegicus	9	1994	bait stations, traps	bromadiolone	Lorvelec & Pascal 2005, Pascal et al. 2005
Trielen, Molène	FRA	R. norvegicus	17	1996	bait stations, traps	chlorofacinone	Lorvelec & Pascal 2005
Bono, Sept-Île	FRA	R. norvegicus	22	1994	bait stations, traps	bromadiolone	Lorvelec & Pascal 2005, Pascal et al. 2005
Tomé	FRA	R. norvegicus	30	2002	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005
Folaca, Cerbicales	FRA	R. rattus	0.2	2001	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005
Burgaux, Martinique	FRA	R. rattus	0.49	2002	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005
Percé, Martinique	FRA	R. rattus	0.54	1999	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005
Toro, Cerbicales	FRA	R. rattus	0.9	1991	hand broadcast		M. Pascal unpublished data
Poirier, Martinique	FRA	R. rattus	2.1	2002	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005
Hardy, Martinique	FRA	R. rattus	2.6	2002	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005
Mato, New Caledonia	FRA	R. rattus	4	1998	hand broadcast	brodifacoum	E. Bell pers. comm.
Plane, Riou	FRA	R. rattus	15	2005	hand broadcast, traps	bromadiolone	M. Pascal unpublished data
Surprise, New Caledonia	FRA	R. rattus	24	2005	hand broadcast	bromadiolone	F. Courchamp pers. comm.
Lavezzu (+16 islets), Lavezzi	FRA	R. rattus	90	2000	hand broadcast, traps	bromadiolone	Lorvelec & Pascal 2005
St. Paul	FRA	R. rattus	800	1996	aerial broadcast, hand broadcast	brodifacoum	Micol & Jouventin 2002; Lorvelec & Pascal 2005
Kastronisia-1, Sporades	GRE	R. norvegicus	2	2006	bait stations	brodifacoum	J. Fric pers. comm.
Kastronisia-2, Sporades	GRE	R. norvegicus	2.5	2006	bait stations	brodifacoum	J. Fric pers. comm.

Kasidis, Lagofytonisia	GRE	R. rattus	<1	2005	bait stations	brodifacoum	J. Fric pers. comm.
Kastronisia-1, Sporades	GRE	R. rattus	2	2006	bait stations	brodifacoum	J. Fric pers. comm.
Kastronisia-2, Sporades	GRE	R. rattus	2.5	2006	bait stations	brodifacoum	J. Fric pers. comm.
Lachanou, Lagofytonisia	GRE	R. rattus	5	2005	bait stations	brodifacoum	J. Fric pers. comm.
Polemika, Lagofytonisia	GRE	R. rattus	11.2	2005	bait stations	brodifacoum	J. Fric pers. comm.
Flatey Island	ICE	M. musculus	50	1971		warfarin	Peterson 1979
Flatey Island	ICE	R. norvegicus	50	1971		warfarin	Peterson 1979
Sangalaki, Derawen Islands	IND	R. rattus	13.5	2003	bait stations	brodifacoum	G. Meier pers. comm.
Gemino di Terra, Elba	ITA	R. rattus	1.8	2000	bait stations	brodifacoum, bromadiolone	N. Baccetti & P. Sposimo pers. comm.; Perfetti et al. 2001; Genovesi 2005
Gemino di Fuori, Elba	ITA	R. rattus	1.9	2000	bait stations	brodifacoum, bromadiolone	N. Baccetti & P. Sposimo pers. comm.; Perfetti et al. 2001; Genovesi 2005
Isola La Scola	ITA	R. rattus	2	2001	bait stations		Perfetti et al. 2001; Genovesi 2005
dei Topi	ITA	R. rattus	10	2000	bait stations	brodifacoum	Perfetti et al. 2001; Genovesi 2005
d'Ercole	ITA	R. rattus	10	2000	bait stations	brodifacoum	Perfetti et al. 2001; Genovesi 2005
Scoglio La Peraiola	ITA	R. rattus	10	2000	bait stations	brodifacoum	Perfetti et al. 2001; Genovesi 2005
Île aux Sables	MAU	M. musculus	8	1995	bait stations	brodifacoum, bromadiolone	Bell 2002
Île Cocos	MAU	M. musculus	15	1995	bait stations	brodifacoum, bromadiolone	Bell 2002
Flat	MAU	M. musculus	253	1998	hand broadcast, bait stations	brodifacoum, bromadiolone	Bell 2002
Gunner's Quoin	MAU	R. norvegicus	65	1996	hand broadcast	brodifacoum, bromadiolone	Bell 2002
Île Aux Aigrettes	MAU	R. rattus	25	1987	hand broadcast	brodifacoum	Merton 1987
Gabriel	MAU	R. rattus	42	1995	hand broadcast	brodifacoum, bromadiolone	Bell 2002
Flat	MAU	R. rattus	253	1998	hand broadcast, bait stations	brodifacoum, bromadiolone	Bell 2002
Rasa, Gulf of California	MEX	M. musculus	60	1994	bait stations, traps	brodifacoum	J. Ramirez pers. comm.

San Jorge West, Gulf of California	MEX	R. rattus	5	2000	bait stations	cholecalciferol	Donlan et al. 2003
San Jorge East, Gulf of California	MEX	R. rattus	5	2000	bait stations	diphacinone	Donlan et al. 2003
San Jorge Middle, Gulf of California	MEX	R. rattus	14	2000	bait stations	brodifacoum	Donlan et al. 2003
Rasa, Gulf of California	MEX	R. rattus	60	1994	bait stations, traps	brodifacoum	J. Ramirez pers. comm.
San Roque, Baja California	MEX	R. rattus	70	1994	bait stations	brodifacoum, bromethalin	Donlan et al. 2000; Tershy et al. 2002
Papakohatu, Hauraki Gulf	NZL	M. musculus	0.7	1996	bait stations	brodifacoum	Clout & Russell 2006
Whenuakura (Whangamata), Whitianga	NZL	M. musculus	2	1984	bait stations	bromadiolone	Clout & Russell 2006
Motutapu, Marlborough Sounds	NZL	M. musculus	2	1989	bait stations	flocoumafen	Clout & Russell 2006
Moturemu, Kaipara Harbor	NZL	M. musculus	5	1992	bait stations	brodifacoum	Clout & Russell 2006
Allports, Marlborough Sounds	NZL	M. musculus	16	1989	bait stations, traps	brodifacoum	Clout & Russell 2006
Rimariki	NZL	M. musculus	22	1991	bait stations	bromadiolone	Clout & Russell 2006
Motutapere, West Coromandel	NZL	M. musculus	45	1994	aerial broadcast, bait stations	brodifacoum	Clout & Russell 2006
Browns (Motukorea), Hauraki Gulf	NZL	M. musculus	58	1995	aerial broadcast	bromadiolone	Clout & Russell 2006
Mokoia, Lake Rotorua	NZL	M. musculus	133	2003	aerial broadcast	brodifacoum	Clout & Russell 2006
Mou Waho, Lake Wanaka	NZL	M. musculus	140	1995	aerial broadcast	brodifacoum	Clout & Russell 2006
Motuihe, Hauraki	NZL	M. musculus	179	1997	aerial broadcast	brodifacoum	Clout & Russell 2006
Mana	NZL	M. musculus	217	1990	bait stations, aerial broadcast	flocoumafen, brodifacoum	Clout & Russell 2006
Enderby, Auckland	NZL	M. musculus	710	1995	aerial broadcast	brodifacoum	Clout & Russell 2006
Arch, Mokohinau	NZL	R. exulans	1	1991	aerial broadcast, hand broadcast	brodifacoum	Clout & Russell 2006
Flax (Hokoromea), Mokohinau	NZL	R. exulans	1	1991	aerial broadcast, hand broadcast	brodifacoum	Clout & Russell 2006
Lizard, Mokohinau	NZL	R. exulans	1	1978	hand broadcast	bromadiolone	Clout & Russell 2006

Motupapa-Stack C (Mokohinau)	NZL	R. exulans	2	1991	aerial broadcast, hand broadcast	brodifacoum	Clout & Russell 2006
Rurima, Bay of Plenty	NZL	R. exulans	4.5	1984	bait stations	1080, bromadiolone	Clout & Russell 2006
Stacks (B-G,I,J), Mokohinau	NZL	R. exulans	10	1991	aerial broadcast, hand broadcast	brodifacoum	Clout & Russell 2006
Maori Bay, Mokohinau	NZL	R. exulans	11	1991	aerial broadcast, hand broadcast	brodifacoum	Clout & Russell 2006
Whangaokena (East), East Cape	NZL	R. exulans	13	1997			Clout & Russell 2006
Atihau (Trig), Mokohinau	NZL	R. exulans	16	1991	aerial broadcast, hand broadcast	brodifacoum	McFadden 1994; Atkinson & Towns 2001
Korapuki, Mercury	NZL	R. exulans	18	1987	bait stations	bromadiolone	Clout & Russell 2006
Middle Chain, Aldermen	NZL	R. exulans	23	1992	aerial broadcast	brodifacoum	Clout & Russell 2006
Double, Mercury	NZL	R. exulans	27	1989	bait stations, hand broadcast	bromadiolone, flocoumafen	Clout & Russell 2006
Motuopao, Far North	NZL	R. exulans	30	1990	bait stations	brodifacoum	Clout & Russell 2006
Burgess, Mokohinau	NZL	R. exulans	56	1991	aerial broadcast	brodifacoum	Clout & Russell 2006
Motuara, Marlborough Sounds	NZL	R. exulans	59	1991	bait stations	brodifacoum	Clout & Russell 2006
Fanal, Mokohinau	NZL	R. exulans	73	1997	aerial broadcast, hand broadcast	brodifacoum	Clout & Russell 2006
Whakaterepapanui, Rangitoto	NZL	R. exulans	74	1999	aerial broadcast	brodifacoum	Clout & Russell 2006
Mauipae (Coppermine), Hen and Chickens	NZL	R. exulans	80	1997	aerial broadcast	brodifacoum	Clout & Russell 2006
Rarotoka (Centre Island)	NZL	R. exulans	88	1997	aerial broadcast	brodifacoum	Clout & Russell 2006
Stanley, Mercury	NZL	R. exulans	100	1992	aerial broadcast, hand broadcast	brodifacoum	Clout & Russell 2006
Mauiroto (Whatupuke), Hen and Chickens	NZL	R. exulans	102	1993	aerial broadcast	brodifacoum	Clout & Russell 2006
Marotiri (Lady Alice), Hen and Chickens	NZL	R. exulans	120	1994	aerial broadcast	brodifacoum	Clout & Russell 2006
Long, Marlborough Sounds	NZL	R. exulans	142	1998	aerial broadcast	brodifacoum	Clout & Russell 2006

Putauhinu, Stewart Island	NZL	R. exulans	145	1997	aerial broadcast	brodifacoum	McClelland 2002a
Cuvier, Coromandel East	NZL	R. exulans	170	1993	aerial broadcast	brodifacoum	Clout & Russell 2006
Tiritiri Matangi, Hauraki Gulf	NZL	R. exulans	220	1993	aerial broadcast	brodifacoum	Clout & Russell 2006
Red Mercury, Mercury	NZL	R. exulans	225	1992	aerial broadcast, hand broadcast	brodifacoum	Clout & Russell 2006
Inner Chetwode (Nukuwaiata), Marlborough Sounds	NZL	R. exulans	242	1994	aerial broadcast, traps	brodifacoum	Clout & Russell 2006
Mayor (Tuhua), Bay of Plenty	NZL	R. exulans	1277	2000	aerial broadcast	brodifacoum	Clout & Russell 2006
Codfish (Whenua Hou), Stewart Island	NZL	R. exulans	1396	1998	aerial broadcast	brodifacoum	Clout & Russell 2006
Kapiti	NZL	R. exulans	1965	1996	aerial broadcast	brodifacoum	Clout & Russell 2006
Raoul, Kermadecs	NZL	R. exulans	2938	2002	aerial broadcast	brodifacoum	Clout & Russell 2006
Hauturu (Little Barrier Island)	NZL	R. exulans	3083	2004	aerial broadcast	brodifacoum	R. Griffiths pers. comm.
Maukaha Rocks (Whangamata)	NZL	R. norvegicus	0.1	1984	bait stations	bromadiolone	Clout & Russell 2006
Black Rocks (17 islets), Bay of Islands	NZL	R. norvegicus	1	1992	aerial broadcast, hand broadcast, bait stations	brodifacoum	Clout & Russell 2006
Maria, Hauraki Gulf	NZL	R. norvegicus	1	1960	bait stations	warfarin	Clout & Russell 2006
Motiti	NZL	R. norvegicus	1	1990		warfarin	Clout & Russell 2006
Motutapu, Bay of Islands	NZL	R. norvegicus	1	1990			Clout & Russell 2006
Motuterakihi	NZL	R. norvegicus	1	1985			Clout & Russell 2006
Taranaki, Bay of Islands	NZL	R. norvegicus	1	1990			Clout & Russell 2006
Tahoramaurea, Kapiti	NZL	R. norvegicus	1	1996	bait stations	brodifacoum	Empson & Miskelly 1999; Clout & Russell 2006
David Rocks, Hauraki Gulf	NZL	R. norvegicus	1	1964	bait stations	warfarin	Moors 1985; Thomas & Taylor 2002, Clout & Russell 2006
Unnamed N. of Cape Wīwiki A (Snail Rock)	NZL	R. norvegicus	1.5	1999			R. Parrish pers. comm.
Whenuakura (Whangamata), Whitianga	NZL	R. norvegicus	2	1984	bait stations	bromadiolone	Clout & Russell 2006

Wainui, Bay of Islands	NZL	R. norvegicus	2	1991			Clout & Russell 2006
Motungara, Kapiti	NZL	R. norvegicus	3	1996	bait stations	brodifacoum	Clout & Russell 2006
Tarahiki, Hauraki Gulf	NZL	R. norvegicus	5	2000	hand broadcast	brodifacoum	Clout & Russell 2006
Moturemu, Kaipara Harbor	NZL	R. norvegicus	5	1992	bait stations	brodifacoum	I. McFadden pers. comm.
Moturemu, Kaipara Harbor	NZL	R. norvegicus	5	2004	bait stations, traps	brodifacoum	Russell et. al. 2005, I. McFadden pers. comm.
Te Haupa (Saddle), Hauraki Gulf	NZL	R. norvegicus	6	1989	hand broadcast, traps	brodifacoum	Veitch & Bell 1990
Hawea, Fiordland	NZL	R. norvegicus	9	1986	bait stations	brodifacoum	Clout & Russell 2006
Motuhoropapa, Hauraki	NZL	R. norvegicus	9.5	1997	bait stations	brodifacoum	Cameron 1998; Clout & Russell 2006
Motuhoropapa, Hauraki	NZL	R. norvegicus	9.5	1991	hand broadcast, traps	bromadiolone, flocoumafen	Cameron 1998; Clout & Russell 2006
Motuhoropapa, Hauraki	NZL	R. norvegicus	9.5	2001	bait stations	brodifacoum	Clout & Russell 2006
Motuhoropapa, Hauraki	NZL	R. norvegicus	9.5	2002	hand broadcast, bait stations	brodifacoum	Clout & Russell 2006
Motuhoropapa, Hauraki	NZL	R. norvegicus	9.5	1984	bait stations	1080, brodifacoum	Moors 1985; Thomas & Taylor 2002; Clout & Russell 2006
Motuhoropapa, Hauraki	NZL	R. norvegicus	9.5	1981	bait stations	1080, brodifacoum	Moors 1985; Clout & Russell 2006
Hauturu, Whangamata	NZL	R. norvegicus	10	1993	bait stations	brodifacoum	Clout & Russell 2006
Patiti (Banded), Lake Rotomahana	NZL	R. norvegicus	12.8	2004	bait stations	brodifacoum	Bancroft 2004
Motu-O-Kura, Hawkes Bay	NZL	R. norvegicus	14	1990	hand broadcast	brodifacoum	Clout & Russell 2006
Otata, Hauraki Gulf	NZL	R. norvegicus	22	2001	bait stations	brodifacoum	Clout & Russell 2006
Otata, Hauraki Gulf	NZL	R. norvegicus	22	2002	hand broadcast, bait stations	brodifacoum	Clout & Russell 2006
Otata, Hauraki Gulf	NZL	R. norvegicus	22	1991	hand broadcast	bromadiolone, flocoumafen	Clout & Russell 2006
Otata, Hauraki Gulf	NZL	R. norvegicus	22	1981	hand broadcast	bromadiolone, flocoumafen	Moors 1985
Otata, Hauraki Gulf	NZL	R. norvegicus	22	1979	bait stations, hand broadcast, traps	1080, brodifacoum	Moors 1985; Veitch & Bell 1990
Pakatoa, Hauraki Gulf	NZL	R. norvegicus	29	1998	bait stations	brodifacoum	Clout & Russell 2006
Titi, Marlborough Sounds	NZL	R. norvegicus	32	1983	hand broadcast	warfarin	Clout & Russell 2006

Matakohe, Limestone	NZL	R. norvegicus	37	1998	bait stations	brodifacoum	Clout & Russell 2006
Browns (Motukorea), Hauraki Gulf	NZL	R. norvegicus	58	1995	aerial broadcast	bromadiolone	Clout & Russell 2006
Puangiangi, Rangitoto	NZL	R. norvegicus	69	2001	aerial broadcast	brodifacoum	Clout & Russell 2006
Whakaterepapanui, Rangitoto	NZL	R. norvegicus	74	1999	aerial broadcast	brodifacoum	Clout & Russell 2006
Quail (Otomahua), Lyttleton Harbour	NZL	R. norvegicus	88	2002	bait stations	brodifacoum	Kavermen et al. 2003
Tinui, Rangitoto	NZL	R. norvegicus	95	1999	aerial broadcast	brodifacoum	Clout & Russell 2006
Mokoia, Lake Rotorua	NZL	R. norvegicus	133	1989	bait stations	brodifacoum, flocoumafen	Clout & Russell 2006
Rakino, Hauraki Gulf	NZL	R. norvegicus	148	2002	bait stations	brodifacoum	Clout & Russell 2006
Moturoa, Bay of Islands	NZL	R. norvegicus	157	2003	hand broadcast, bait stations, traps	brodifacoum	Clout & Russell 2006
Moturoa, Bay of Islands	NZL	R. norvegicus	157	1993	hand broadcast, bait stations, traps	brodifacoum	Clout & Russell 2006
Breaksea, Fiordland	NZL	R. norvegicus	170	1988	bait stations	brodifacoum	Taylor & Thomas 1993; Clout & Russell 2006
Moutohora (Whale), Bay of Plenty	NZL	R. norvegicus	173	1986	aerial broadcast	brodifacoum	Clout & Russell 2006
Motuihe, Hauraki	NZL	R. norvegicus	179	1997	aerial broadcast	brodifacoum	Clout & Russell 2006
Ulva, Stewart Island	NZL	R. norvegicus	267	1996	bait stations, traps	brodifacoum, bromadiolone	Thomas & Taylor 2002; Clout and Russell 2006
Mayor (Tuhua), Bay of Plenty	NZL	R. norvegicus	1277	2000	aerial broadcast	brodifacoum	Clout & Russell 2006
Kapiti	NZL	R. norvegicus	1965	1996	aerial broadcast	brodifacoum	Clout & Russell 2006
Raoul, Kermadecs	NZL	R. norvegicus	2938	2002	aerial broadcast	brodifacoum	Clout & Russell 2006; M. Abrose pers. comm.
Campbell	NZL	R. norvegicus	11300	2002	aerial broadcast	brodifacoum	Clout & Russell 2006
Koi, Hauraki Gulf	NZL	R. rattus	0.28	1997	hand broadcast, traps	brodifacoum	Lee 1999

Black Rocks (17 islets), Bay of Islands	NZL	R. rattus	1	1992	aerial broadcast, hand broadcast, bait stations	brodifacoum	Clout & Russell 2006
Mokopuna (Leper), Wellington Harbor	NZL	R. rattus	1	1990			Clout & Russell 2006
Unnamed N. of Cape Wiwiki A (Snail Rock)	NZL	R. rattus	1.5	1994			R. Parrish pers. comm.
Awaiti, Marlborough Sounds	NZL	R. rattus	2	1982	bait stations	brodifacoum	Clout & Russell 2006
Duffers Reef, Marlborough Sounds	NZL	R. rattus	2	1983	bait stations	brodifacoum	Veitch & Bell 1990
Haulashore, Nelson	NZL	R. rattus	6	1991	bait stations	brodifacoum	Clout & Russell 2006
Iona, Stewart	NZL	R. rattus	7	2004	bait stations	brodifacoum	Clout & Russell 2006
Hokianga, Ohiwa Harbour	NZL	R. rattus	8	2005	bait stations	brodifacoum	D. Paine pers. comm.
Goat	NZL	R. rattus	10	1992	bait stations	brodifacoum	Clout & Russell 2006
Goat	NZL	R. rattus	10	2005	traps, bait stations (but no uptake)	brodifacoum	Clout & Russell 2006
Harakeke , Bay of Islands	NZL	R. rattus	12	1994			Veitch 1995; D. P. Taylor pers. comm.
Patiti (Banded), Lake Rotomahana	NZL	R. rattus	12.8	2004	bait stations	brodifacoum	Bancroft 2004
Tawhitinui	NZL	R. rattus	21	1983	bait stations	brodifacoum	Clout & Russell 2006
Somes (Matiu), Wellington Harbor	NZL	R. rattus	32	1988		brodifacoum	Clout & Russell 2006
Motutapere, West Coromandel	NZL	R. rattus	45	1994	aerial broadcast, bait stations	brodifacoum	Clout & Russell 2006
Motutapere, West Coromandel	NZL	R. rattus	46	2005	aerial broadcast	brodifacoum	J. Russell unpublished data
Moturoa, Bay of Islands	NZL	R. rattus	157	1993	hand broadcast, bait stations, traps	brodifacoum	Clout & Russell 2006
Selvagem Grande, Madiera	POR	M. musculus	200	2003	bait stations, hand broadcast	brodifacoum	J. Parkes unpublished data
Frégate, Seychelles	ROS	M. musculus	219	2000	aerial broadcast	brodifacoum	Merton et al. 2002
Frégate, Seychelles	ROS	R. norvegicus	219	2000	aerial broadcast	brodifacoum	Merton et al. 2002

Bird, Seychelles	ROS	R. rattus	101	1996	hand broadcast, bait stations	brodifacoum	Merton et al 2002; B. Simmons pers. comm.
Denis, Inner Seychelles	ROS	R. rattus	143	2003		brodifacoum	D. Merton pers. comm.
Denis, Inner Seychelles	ROS	R. rattus	143	2000	aerial broadcast	brodifacoum	Merton et al. 2002
Curieuse, Seychelles	ROS	R. rattus	286	2000	aerial broadcast	brodifacoum	Merton et al. 2002
Praslin, East Coast	SL	R. rattus	1.1	1993	bait stations	brodifacoum	J. Daltry pers. comm.
Praslin, East Coast	SL	R. rattus	2.1	2000	bait stations	brodifacoum	J. Daltry pers. comm.
Conills, Balearic	SPA	R. rattus	1	1999			Genovesi 2005
Ray Francisco, Chafarines	SPA	R. rattus	12	1992	bait stations	brodifacoum	Orueta et al. 2005
Ray Francisco, Chafarines	SPA	R. rattus	12	2000	bait stations	flocoumafen, brodifacoum, bromadiolone	Orueta et al. 2005
Chumbe	TAN	R. rattus	20	1997			http://www.earthfoot.org/places/tz001b.htm
Oeno, Pitcairn	UK	R. exulans	62	1997	hand broadcast	brodifacoum	E. Bell pers. comm.
Ducie, Pitcairn	UK	R. exulans	74	1997	hand broadcast	brodifacoum	E. Bell pers. comm.
Calf Islet, Falklands	UK	R. norvegicus	<1	2001	hand broadcast	brodifacoum	Brown et al. 2001
Rat Island, Falklands	UK	R. norvegicus	0.5	2001	hand broadcast	brodifacoum	Brown et al. 2001
Calf Island, Falklands	UK	R. norvegicus	5	2001	hand broadcast	brodifacoum	Brown et al. 2001
Horse, Falklands	UK	R. norvegicus	5	2001	bait stations	brodifacoum	Brown et al. 2001
Nonsuch, Bermuda	UK	R. norvegicus	5.8	<1985	bait stations	warfarin, diphacinone	Wingate 1985
Bottom Tussac, Falklands	UK	R. norvegicus	8	2001	bait stations	brodifacoum	Brown et al. 2001
Double, Falklands	UK	R. norvegicus	9	2001	hand broadcast	brodifacoum	Brown et al. 2001
Top Tussac, Falklands	UK	R. norvegicus	12	2001	bait stations	brodifacoum	Brown et al. 2001
Cardigan	UK	R. norvegicus	16	1980			Johnstone et al. 2005
Outer, Falklands	UK	R. norvegicus	20	2001	hand broadcast	brodifacoum	Brown et al. 2001
Puffin (Seiriol's Island), Isle of Anglesey	UK	R. norvegicus	32	1998			Johnstone et al. 2005, J. Hughes pers. comm.
Ailsa Craig	UK	R. norvegicus	104	1991	bait stations, hand broadcast	warfarin	Zonfrillo 2002; Geovesi 2005, B. Zonfrillo pers. comm.

Ramsey, Pembrokeshires	UK	R. norvegicus	253	2000	hand broadcast	difenacoum	J. Hughes pers. comm., E. Bell pers. comm.
Handa	UK	R. norvegicus	363	1997			J. Hughes pers. comm.
Lundy, Bristol Channel	UK	R. norvegicus	430	2004	bait stations	difenacoum, bromadiolone	E. Bell pers. comm.; K. Varnham pers. comm.
Harpoon, Falklands	UK	R. norvegicus		2001	hand broadcast	brodifacoum	Brown et al. 2001
William Dean Cay, Turks & Caicos	UK	R. rattus	0.6	2002	bait stations	brodifacoum	G. Gerber pers. comm.
Pusey Cay, Turks & Caicos	UK	R. rattus	0.7	2002	bait stations	brodifacoum	G. Gerber pers. comm.
Sim Cay, Turks & Caicos	UK	R. rattus	0.8	2002	bait stations	brodifacoum	G. Gerber pers. comm.
Bay Cay, Turks & Caicos	UK	R. rattus	3.6	2002	bait stations	brodifacoum	G. Gerber pers. comm.
Sandy Cay (White Cay), British Virgin	UK	R. rattus	5.7	2002	bait stations	brodifacoum	Varnham 2003
Nonsuch, Bermuda ^b	UK	R. rattus	5.8	2005	bait stations		Maderios 2005
Nonsuch, Bermuda	UK	R. rattus	5.8	<1985	bait stations	warfarin, diphacinone	Wingate 1985
Lundy, Bristol Channel	UK	R. rattus	430	2004	bait stations	difenacoum, bromadiolone	E. Bell pers. comm.; K. Varnham pers. comm.
Rose Atoll, American Samoa	USA	R. exulans	6		poison, traps	brodifacoum, bromethalin	K. Swift pers. comm.
Kure Atoll, Hawaii	USA	R. exulans	105	1993	bait stations, traps	brodifacoum, bromethalin	K. Swift pers. comm.
Stephen Cay, US Virgin	USA	R. rattus	0.8	<1992			Garcia et al. 2002
Midway Atoll-Spit, Hawaii	USA	R. rattus	0.8	1994	traps		K. Swift pers. comm.
Mokoli'i, Hawaii	USA	R. rattus	1.5	2002	bait stations, traps	diphacinone	Smith et al. 2006
Green Cay, US Virgin	USA	R. rattus	14	2000	traps		M. Evans pers. comm.
Monito, Puerto Rico	USA	R. rattus	15	1999	bait stations, hand broadcast	brodifacoum	Garcia et al. 2002
Anacapa East, Channel Islands	USA	R. rattus	43	2001	aerial broadcast, hand broadcast	brodifacoum	Howald et al. 2005
Anacapa Middle, Channel Islands	USA	R. rattus	71	2002	aerial broadcast, hand broadcast	brodifacoum	Howald et al. 2005
Buck, US Virgin	USA	R. rattus	71	2000	bait stations	diphacinone	Witmer et al. 2007

Midway Atoll-Eastern, Hawaii	USA	R. rattus	133	1995	bait stations, traps	brodifacoum, bromethalin	K. Swift pers. comm.
Anacapa West, Channel Islands	USA	R. rattus	182	2002	aerial broadcast, hand broadcast	brodifacoum	Howald et al. 2005
Midway Atoll-Sand, Hawaii	USA	R. rattus	486	1996	bait stations, traps	brodifacoum, bromethalin	K. Swift pers. comm.
Cayo Ratones, Puerto Rico	USA	R. rattus		<1992			Garcia et al. 2002

FAILED ERADICATIONS OR ERADICATIONS OF UNKNOWN OUTCOME

<i>Island (Alternative Name), Group</i>	<i>County^a</i>	<i>Species</i>	<i>Area (ha)</i>	<i>Year Eradicated</i>	<i>Methods (1°, 2°, 3°)</i>	<i>Rodenticide (1°, 2°, 3°)</i>	<i>Status</i>	<i>Reference</i>
West Maiden, West Coast	A&B	R. rattus	2	1998	bait stations	brodifacoum	unknown	J. Daltry pers. comm.
Codrington, North Sound	A&B	R. rattus	8	1998	bait stations	brodifacoum	unknown	J. Daltry pers. comm.
Maiden (2 islets), North Sound	A&B	R. rattus	9	1999	bait stations	brodifacoum	unknown	J. Daltry pers. comm.
Varanus, Lowendal	AUS	M. musculus	83	1993	bait stations	1080	failed	Burbidge & Morris 2002
Adele	AUS	R. exulans	217	2004			unknown	A. Burbidge unpublished data
Shelter	AUS	R. rattus	8	2003		brodifacoum	unknown	A. Burbidge unpublished data
Delta, Montebellos	AUS	R. rattus	38	1996	bait stations	brodifacoum	failed	Burbidge 2004
Crocus, Montebellos	AUS	R. rattus	41	1996	bait stations	brodifacoum	failed	Burbidge 2004
Primose, Montebellos	AUS	R. rattus	41	1996	bait stations	brodifacoum	failed	Burbidge 2004
Campbell, Montebellos	AUS	R. rattus	47	1996	bait stations	brodifacoum	failed	Burbidge 2004
Hermite, Montebellos	AUS	R. rattus	1022	1996	bait stations	brodifacoum	failed	Burbidge 2004
Hermite, Montebellos	AUS	R. rattus	1022	1999	aerial	brodifacoum	failed	Burbidge 2004
Low Cay, San Salvador	BAH	R. rattus	10.8	1999	bait stations	brodifacoum	failed	Hayes et al. 2004
Motu To'u, Suvarrow	COI	R. exulans	10	2003	hand broadcast	brodifacoum	failed	G. Wragg pers. comm
Marielas Grande, Galapagos	ECU	R. rattus	1.3	2004	bait stations	brodifacoum	unknown	K. Campbell pers. comm.
Bainbridge-1, Galapagos	ECU	R. rattus	11.4	2002	hand broadcast	brodifacoum	failed	K. Campbell pers. comm.
Pinzon, Galapagos	ECU	R. rattus	1815	1987	bait stations, hand broadcast	coumatetralyl, brodifacoum, 1080	failed	K. Campbell pers. comm.
Île du Château, Kerguelen	FRA	M. musculus	250			brodifacoum	unknown	J. Chapuis pers. comm., M Pascal pers. comm.
Motu-o-ari, Gambier Group, French Polynesia	FRA	R. exulans	3	2003	hand broadcast	brodifacoum	unknown	G. Wragg pers. comm

Signal, New Caledonia	FRA	R. exulans	6	1998	hand broadcast	brodifacoum	failed	E. Bell pers. comm.
Teuaua/Ua-Uka, Marquesas, French Polynesia	FRA	R. exulans	7	1988		brodifacoum, chlorofacinone	failed	Sechan 1987
Makapu, Gambier Group, French Polynesia	FRA	R. exulans	12	2003	hand broadcast	brodifacoum	unknown	G. Wragg pers. comm
Mekiro, Gambier Group, French Polynesia	FRA	R. exulans	15	2003	hand broadcast	brodifacoum	unknown	G. Wragg pers. comm
Teanaone & Tepapuri, Gambier Group, French Polynesia	FRA	R. exulans	30	2003	hand broadcast	brodifacoum	unknown	G. Wragg pers. comm
Vahanga, Tuamotu Group, French Polynesia	FRA	R. exulans	382	2000		brodifacoum, chlorofacinone	failed	M. Pascal unpublished data
6 islets, St.Riom	FRA	R. norvegicus	1.7	2000	hand broadcast. traps	bromadiolone	failed	Lorvelec & Pascal 2005
St. Riom , St. Riom	FRA	R. norvegicus	14	2000	hand broadcast, traps	bromadiolone	failed	Lorvelec & Pascal 2005
Glénanm Le Loc'h	FRA	R. norvegicus		2003	hand broadcast, traps	bromadiolone	failed	M. Pascal unpublished data
Folaccheda, Cerbicales	FRA	R. rattus	0.1	2001	hand broadcast. traps	bromadiolone	unknown	Lorvelec & Pascal 2005
Teuaua/Ua-Uka, Marquesas, French Polynesia	FRA	R. rattus	7	1987		brodifacoum, chlorofacinone	failed	Sechan 1987
Fajou, Guadeloupe	FRA	R. rattus	120	2002	hand broadcast, traps	bromadiolone	failed	Lorvelec & Pascal 2005, Abdelkrim et al. 2005b
Île du Château, Kerguelen	FRA	R. rattus	250			brodifacoum	unknown	J. Chapuis pers. comm., M. Pascal pers. comm.
Isabel	MEX	R. rattus	194	1995	bait stations	brodifacoum, hand broadcast	failed	Rodriguez et al. 2006
Te Haupa (Saddle), Hauraki Gulf	NZL	M. musculus	6	1993	bait stations	flocoumafen	failed	Clout & Russell 2006
Patiti (Banded), Lake Rotomahana	NZL	M. musculus	12.8	2004	bait stations	brodifacoum	failed	Bancroft 2004
Matakohe (Limestone/Whangarei)	NZL	M. musculus	37	1998	aerial	brodifacoum	failed	Clout & Russell 2006
Matakohe (Limestone/Whangarei)	NZL	M. musculus	37	1997	aerial	brodifacoum	failed	Clout & Russell 2006
Matakohe (Limestone/Whangarei)	NZL	M. musculus	37	2001	bait stations	brodifacoum	failed	Clout & Russell 2006
Mokoia, Lake Roturua	NZL	M. musculus	133	1996	aerial	brodifacoum	failed	Clout & Russell 2006

Coppermine (Mauipae), Hen and Chickens	NZL	R. exulans	80	1992	bait stations	brodifacoum	failed	Thomas & Taylor 2002; Towns & Broome 2003
Motuhoropapa, Hauraki	NZL	R. norvegicus	9.5	1978	traps		failed	Thomas & Taylor 2002
Rotoroa, Hauraki Gulf	NZL	R. norvegicus	90	1992	bait stations	brodifacoum	failed	Lee 1999; Clout & Russell 2006
Rakino, Hauraki Gulf	NZL	R. norvegicus	148	1992	bait stations	brodifacoum	failed	Clout & Russell 2006
Moturako, Great Barrier Island	NZL	R. rattus	1	1990	hand broadcast	flocoumafen	unknown	G. Taylor pers. comm.
Oyster, Great Barrier Island	NZL	R. rattus	1	1990	hand broadcast	flocoumafen	unknown	G. Taylor pers. comm.
Wood Stack A, Great Barrier Island	NZL	R. rattus	1	1990	hand broadcast	flocoumafen	unknown	G. Taylor pers. comm.
Wood, Great Barrier Island	NZL	R. rattus	1	1990	hand broadcast	flocoumafen	unknown	G. Taylor pers. comm.
Saddle, Great Barrier Island	NZL	R. rattus	2	1990	hand broadcast	flocoumafen	unknown	G. Taylor pers. comm.
Opakau, Great Barrier Island	NZL	R. rattus	4	1990	hand broadcast	flocoumafen	unknown	G. Taylor pers. comm.
Frégate, Seychelles	ROS	R. norvegicus	219	1996	bait stations	brodifacoum	failed	Thorsen et al. 2000
North, Seychelles	ROS	R. rattus	210	2003	aerial broadcast	brodifacoum	failed	D. Merton, pers. comm.
Pitcairn, Pitcairn	UK	R. exulans	500	1998	hand broadcast	brodifacoum	failed	E. Bell pers. comm.
Monito, Puerto Rico	USA	R. rattus	15	1993	hand broadcast	brodifacoum	failed	Garcia et al. 2002
Palmyra, Line Islands	USA	R. rattus	230	2001	bait stations	brodifacoum	failed	G. Howald unpublished data

^a Country abbreviations: A&B, Antigua & Barbuda; AUS, Australia; BAH, Bahamas; CAN, Canada; COI, Cook Islands; ECU, Ecuador; FRA, France; ICE, Iceland; IND, Indonesia; ITA, Italy; MAU, Mauritius; MEX, Mexico; NZL, New Zealand; ROI, Republic of Seychelles; SL, Saint Lucia; SPA, Spain; TAN, Tanzania; UK, United Kingdom, USA, United States

^b small area eradicated during initial invasion

LITERATURE CITED

Abbott, I., and A. A. Burbidge. 1995. The occurrence of mammal species on islands of Australia. CALMScience **1**:259-324.

Abdelkrim, J., M. Pascal, C. Calmet, and S. Samadi. 2005a. Importance of assessing population genetic structure before eradication of invasive species: examples from insular Norway Rat populations. Conservation Biology **19**:1509-1518.

- Abdelkrim, J., M. Pascal, and S. Samadi. 2005b. Island colonization and founder effects: the invasion of Guadeloupe islands by ship rats (*Rattus rattus*). *Molecular Ecology* **14**:2923-2931.
- Atkinson, I. A. E., and D. R. Towns. 2001. Advances in New Zealand mammalogy 1990-2000: Pacific rat. *Journal of the Royal Society of New Zealand* **31**:99-109.
- Bancroft, B. 2004. Patiti (Banded) Island rodent eradication 2004. Waimangu Volcanic Valley Ltd and the Department of Conservation, Rotorua, New Zealand.
- Bell, B. D. 2002. The eradication of alien mammals from five offshore islands, Mauritius, Indian Ocean. Pages 40-45 in C. R. Veitch, and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. IUCN SSC Invasive Species Specialist Group, Gland.
- Brown, D., L. Chadderton, and K. Brown. 2001. Report on the Falklands Conservation Rat Eradication Project. http://www.falklandsconservation.com/wildlife/conservation_issues/rat_eradication-final.html#An2.
- Buck, P. E. F. 1995. Field test of rata+ weather block on Lucy Island, British Columbia, Canada. Page 13. Husta Forestry Services LTD., Masset.
- Burbidge, A. A. 2004. Montebello Renewal: Western Shield review—February 2003. *Conservation Science Western Australia* **5**:194-201.
- Burbidge, A. A., and K. D. Morris. 2002. Introduced mammal eradications for nature conservation on western Australian islands: a review. Pages 64-70 in C. R. Veitch, and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. IUCN SSC Invasive Species Specialist Group, Gland.
- Cameron, E. 1998. Bot Soc trips to the Noises (Hauraki Gulf) and an updated species list. *Auckland Botanical Society Journal* **53**:23-25.
- Clout, M. N., and J. C. Russell. 2006. The eradication of introduced mammals from New Zealand Islands. Pages 127-141 in F. Koike, M. N. Clout, M. Kawamichi, M. De Poorter, and K. Iwatsuki, editors. *Assessment and Control of Biological Invasion Risks*. IUCN, Gland.
- Donlan, C. J., G. R. Howald, B. R. Tershy, and D. A. Croll. 2003. Evaluating alternative rodenticides for island conservation: roof rat eradication from the San Jorge Islands, Mexico. *Biological Conservation* **14**:29-34.

- Donlan, C. J., B. R. Tershy, B. S. Keitt, B. Wood, J. A. Sanchez, A. Weinstein, D. A. Croll, and J. L. Alguilar. 2000. Island conservation action in northwest Mexico. Pages 330-338 in D. H. Browne, H. Chaney, and K. Mitchell, editors. Proceedings of the Fifth California Islands Symposium. Santa Barbara Museum of Natural History, Santa Barbara, California, USA.
- Empson, R. A., and C. M. Miskelly. 1999. The risks, costs and benefits of using brodifacoum to eradicate rats from Kapiti Island, New Zealand. *New Zealand Journal of Ecology* **23**:241-254.
- Garcia, M. A., C. E. Diez, and A. O. Alvarez. 2002. The eradication of *Rattus rattus* from Monito Island, West Indies. Pages 116-119 in C. R. Veitch, and M. N. Clout, editors. Turning the tide: the eradication of invasive species. IUCN SSC Invasive Species Specialist Group, Gland.
- Genovesi, P. 2005. Eradications of invasive alien species in Europe: a review. *Biological Invasions* **7**:127-133.
- Hayes, W. K., R. L. Carter, S. C. Jr., and B. Thonton. 2004. Conservation of an Endangered Bahamian Rock Iguana, I. Pages 232-257 in A. C. Alberts, R. L. Carter, W. K. Hayes, and E. P. Martins, editors. Iguanas: Biology and Conservation. University of California, Berkeley.
- Howald, G. R., K. R. Faulkner, B. Tershy, B. Keitt, H. Gellerman, E. M. Creel, M. Grinnel, S. Ortega, and D. A. Croll. 2005. Eradication of black rat from Anacapa Island: biological and social considerations. Pages 299-312 in D. K. Garcelon, and C. A. Schwemm, editors. Proceedings of the Sixth California Islands Symposium. Institute for Wildlife Studies, Arcata, California.
- Johnstone, I. G., C. M. Gray, and D. G. Noble 2005. The state of birds in Wales 2004. RSPB Cymru, Cardiff, United Kingdom.
- Kaiser, G. W., R. H. Taylor, P. D. Buck, J. E. Elliot, G. R. Howald, and M. C. Dever. 1997. The Langara islands seabird habitat recovery project: eradication of Norway rats - 1993-1997 Technical Report Series No. 304. Canadian Wildlife Service, Pacific and Yukon Region, British Columbia.
- Kavermann, M., M. Bowie, and A. Paterson 2003. The eradication of mammalian predators from Quail Island, Banks Peninsula, Canterbury, New Zealand. Lincoln University Wildlife Management Report No. 29. Department of Conservation and the Quail Island Ecological Restoration Trust, Christchurch, New Zealand.
- Lee, M. 1999. Biota of seven islets off Waiheke Island, Inner Hauraki Gulf. *Tane* **37**:99-136.
- Lorvelec, O., and M. Pascal. 2005. French attempts to eradicate non-indigenous mammals and their consequences for native biota. *Biological Invasions* **7**:135-140.

- Maderios, J. 2005. Report on the 2004-2005 Cahow nesting season: continued progress and recovery. Bermuda Audubon Society Newsletter **16**.
- McClelland, P. 2002. Eradication of Pacific rats (*Rattus exulans*) from Whenua Hou Nature Reserve (Codfish Island), Putauhinu and Rarotoka Islands, New Zealand. Pages 173-181 in C. R. Veitch, and M. N. Clout, editors. Turning the tide: the eradication of invasive species. IUCN SSC Invasive Species Specialist Group, Gland.
- McFadden, I., and T. Greene 1994. Using brodifacoum to eradicate kiore (*Rattus exulans*) from Burgess Island and the Knights Group of the Mokohinau Islands. Department of Conservation, Science and Research Series No. 70, Wellington, New Zealand.
- Merton, D. 1987. Eradication of rabbits from Round Island, Mauritius: A conservation success story. Dodo **24**:19-43.
- Merton, D., G. Climo, V. Laboudallon, S. Robert, and C. Mander. 2002. Alien mammal eradication and quarantine on inhabited islands in the Seychelles. Pages 182-198 in C. R. Veitch, and M. N. Clout, editors. Turning the tide: the eradication of invasive species. IUCN SSC Invasive Species Specialist Group, Gland.
- Micol, T., and P. Jouventin. 2002. Eradication of rats and rabbits from Saint-Paul Island, French Southern Territories. Pages 199-205 in C. R. Veitch, and M. N. Clout, editors. Turning the tide: the eradication of invasive species. IUCN SSC Invasive Species Specialist Group, Gland.
- Moors, P. J. 1985. Conservation of island birds: case studies for the management of threatened island birds. International Council for Bird Preservation Technical Publication No. 3.
- Morris, K. D. 1989. Feral animal control on Western Australian islands. Pages 105-111 in A. Burbidge, editor. Australian and New Zealand islands: nature conservation values and management. Department of Conservation and Land Management, Perth.
- Morris, K. D. 2002. The eradication of the black rat (*Rattus rattus*) on Barrow and adjacent islands off the northwest coast of Western Australia. Pages 219-225 in C. R. Veitch, and M. N. Clout, editors. Turning the tide: the eradication of invasive species. IUCN SSC Invasive Species Specialist Group. IUCN, Gland, Switzerland.
- Orueta, J. F., Y. Arranda, T. Gomez, G. G. Tapia, and L. Sanchez-Marmol. 2005. Successful eradication of invasive rodents from a small island through pulsed baiting inside covered stations. Biological Invasions **7**:141-147.
- Pascal, M., F. Siorat, O. Lorvelec, P. Yésou, and D. Simberloff. 2005. A pleasing Norway rat eradication consequence: two shrew species recover. Diversity and Distribution **11**:193-198.

- Perfetti, A., P. Sposimo, and N. Baccetti. 2001. Il controllo dei ratti per la conservazione degli uccelli marini nidificanti nelle isole italiane e mediterranee. *Avocetta* **25**:126.
- Petersen, A. 1979. The breeding birds of Flatey and some adjoining islets, in Breidafjordur, NW Iceland. *Natturufraedingurinn* **49**:229-256.
- Rodríguez, C., R. Torres, and H. Drummond. 2006. Eradicating introduced mammals from a forested tropical island. *Biological Conservation* **130**:98-105.
- Russell, J. C., and M. N. Clout. 2005. Rodent incursions on New Zealand islands in J. Parkes, M. Statham, and G. Edwards, editors. *Proceedings of the 13th Australasian Vertebrate Pest Conference*. Landcare Research, Lincoln, New Zealand.
- Sechan, Y. 1987. Lutte contre les rats en Polynésie française. Dératisation de l'îlot TEUAUA, UA-UKA, Archipel des Marquises. Rapport ORSTOM ref: 09/87/ITRM/DOC-ENT: 0-37.
- Smith, D. G., E. K. Shiinoki, and E. A. VanderWerf. 2006. Recovery of native species following rat eradication on Mokoli'i, O'ahu, Hawai'i. *Pacific Science* **60**:299-303.
- Taylor, R. H., and B. W. Thomas. 1993. Rats eradicated from rugged Breaksea island (170 Ha), Fiordland, New Zealand. *Biological Conservation* **65**:191-198.
- Tershy, B. R., C. J. Donlan, B. Keitt, D. Croll, J. A. Sanchez, B. Wood, M. A. Hermosillo, and G. Howald. 2002. Island conservation in northwest Mexico: A conservation model integrating research, education and exotic mammal eradication. Pages 293-300 in C. R. Veitch, and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. IUCN SSC Invasive Species Specialist Group. IUCN, Gland, Switzerland.
- Thomas, B. W., and R. H. Taylor. 2002. A history of ground-based rodent eradication techniques developed in New Zealand, 1959-1993. Pages 301-310 in C. R. Veitch, and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. IUCN SSC Invasive Species Specialist Group, Gland.
- Thorsen, M., R. Shorten, R. Lucking, and V. Lucking. 2000. Norway rats (*Rattus norvegicus*) on Frégate Island, Seychelles: the invasion; subsequent eradication attempts and implications for island's fauna. *Biological Conservation* **96**:133-138.
- Varnham, K. 2003. Eradication of black rats (*Rattus rattus*) from Sandy Cay, British Virgin Islands. Island Resources Foundation, Tortola, British Virgin Islands.

- Veitch, C. R. 1995. Habitat repair: a necessary prerequisite to translocation of threatened birds. Pages 97-104 in M. Serena, editor. Reintroduction Biology of Australian and New Zealand Fauna. Surrey Beatty & Sons, Chipping Norton.
- Veitch, C. R., and B. D. Bell. 1990. Eradication of introduced animals from the islands of New Zealand. Pages 137-146 in D. R. Towns, C. H. Daugherty, and I. A. E. Atkinson, editors. Ecological restoration of New Zealand islands. Department of Conservation, Wellington, New Zealand.
- Wingate, D. B. 1985. The restoration of Nonsuch Island as a living museum of Bermuda's pre-colonial terrestrial biome. Pages 225-238 in P. J. Moors, editor. Conservation of island birds. International Council for Bird Preservation, Cambridge.
- Witmer, G. W., F. Boyd, and Z. Hillis-Star. 2007. The successful eradication of introduced roof rats (*Rattus rattus*) on Buck Island using diphacinone, followed by resource recovery and an irruption of house mice (*Mus musculus*). Wildlife Research **34**:108-115.
- Zonfrillo, B. 2002. Puffins return to Ailsa Craig. Scottish Bird News **66**:1-2.